

ENZALUTAMIDE-INDUCED HEART FAILURE WITH REDUCED EJECTION FRACTION

*Mirjana Todorović-Mitić¹, Marija Todorović-Marković², Ana Cvetanović^{1,3},
Milan Čevrljaković¹, Ljubomir Dinić^{4,5}*

Prostate cancer is the second most common malignancy in men and the second leading cause of cancer death. Patients unfit for surgery, with locally advanced or metastatic disease, require androgen-deprivation therapy.

In the past decade, enzalutamide and abiraterone were established in prostate cancer treatment. Their safety profile and clinical evidence showed a significant possibility of cardiovascular complications. Heart failure is more common in patients treated with abiraterone. We report a case of a patient with heart failure induced by enzalutamide. In the review of the literature, enzalutamide-induced heart failure was encountered in only two cases.

Further research on the influence of novel anti-androgen therapy on the cardiovascular system, personalized evaluation of the cardiovascular system, and risk stratification for potential cardiovascular adverse events before initiating therapy are needed.

Acta Medica Medianae 2025;64(3): 118–123.

Key words: enzalutamide, heart failure, prostate cancer

¹University Clinical Center Niš, Department of Oncology, Niš, Serbia

²Copenhagen University Hospital Rigshospitalet, Department of Cardiology, Copenhagen, Denmark

³University of Niš, Faculty of Medicine, Department of Oncology, Niš, Serbia

⁴University of Niš, Faculty of Medicine, Department of Surgery and Anesthesiology and Reanimatology, Niš, Serbia

⁵University Clinical Center Niš, Department of Urology, Niš, Serbia

Contact: Mirjana Todorović-Mitić
2a Zorana Radosavljevića Čupe St., 18000 Niš, Serbia
E-mail: mirjanatdrv@gmail.com

diagnosed cases annually, prostate cancer is the second most common malignancy (after lung cancer) and the third leading cause of cancer death (after lung and colorectal cancer) in Serbia (3). Although prostate cancer is highly prevalent, the 5-year relative survival rate for men with localized or locally advanced disease is outstanding, nearing 100%, and yet men with metastatic disease have a 5-year survival rate of only about 30% (4). The risk of developing prostate cancer increases with age, and the median age at diagnosis is 67 years (2).

The etiology and pathogenesis remain unclear but are believed to be multifactorial and include: ageing, ethnicity, hormonal, and genetic factors. Androgens are crucial in the development, maintenance, and progression of prostate cancer. In 80–90% of cases, increased androgen activity is present in the early stage of the disease. The incorporation of androgen deprivation therapies into the treatment strategy for prostate cancer has provided numerous benefits, including enhanced survival for patients with clinically localized or locally advanced disease and improved symptom management for those with advanced stages of the condition (2). First-line androgen deprivation therapy (ADT) mechanism of action is blocking testosterone production by inhibition of the hypothalamic-pituitary-gonadal axis (5). There are different forms of ADT, and the most frequently used ones are gonadotropin-releasing hormone (GnRH) agonists and antagonists. Most metastatic prostate cancer patients, treated with first-line

Introduction

Prostate cancer has one of the highest incidences among malignant diseases, and it has been estimated that more than 11% of males worldwide will have prostate cancer diagnosed during their lifetime. It is also responsible for nearly 50% of cancer deaths (1, 2). It is estimated that 288,300 men will be diagnosed with prostate cancer and 34,700 men will die of the disease by the end of 2025 in the United States of America. With approximately 2,300 to 2,400 newly

ADT, eventually develop castration resistance, as quickly as 7 months after therapy initiation. In that manner, metastatic or non-metastatic hormone-sensitive prostate cancer (HSPC) is becoming castration (hormone) resistant prostate cancer (CRPC), which is associated with a lower survival rate and quality of life deterioration (6). In the past decade, novel hormonal therapy has been introduced as the primary treatment of advanced prostate cancer to overcome castration resistance. Abirateron acetate, enzalutamide, apalutamide and darolutamide are the key novel hormonal therapies that inhibit androgen receptor (AR) function by blocking its dimerization, nuclear translocation, and binding AR-dimers to AR and have been shown to improve overall survival when added to first-line ADT (7).

Cardiac adverse drug reactions and mortality incidence are higher in patients treated with GnRH analogs, especially in individuals with pre-existing cardiac conditions. Incorporating novel agents, such as enzalutamide and abiraterone, into the treatment regimen further elevates cardiovascular risk (8). Abiraterone and enzalutamide are associated with a higher risk of hypertension, and abiraterone is also linked to an increased incidence of heart failure (9). Prescribing information for abiraterone cites risks of cardiac arrhythmias, chest discomfort, and heart failure, while enzalutamide cites only cardiac ischemia risk (10, 11).

Enzalutamide acts as a direct androgen receptor antagonist with greater affinity than first-generation antagonists. It also inhibits its nuclear translocation, thereby blocking the transcription of oncogenic genes essential for cancer growth and survival (12). It was shown that adding enzalutamide to ADT increases overall survival and quality of life in patients who progressed during ADT (13). The efficacy of enzalutamide was investigated in patients with non-metastatic CRPC (defined as prostate-specific antigen (PSA) doubling time \leq 10 months during ADT) (14).

There is an urgent need for further research on the pharmacovigilance of novel hormonal therapy in prostate cancer patients due to the high mortality rate caused by cardiovascular disease, rather than malignancy itself. A recent study done in 2021 showed a higher incidence of cardiac events in patients treated with abiraterone in contrast to the patients treated with enzalutamide and concluded that enzalutamide has a much better safety profile than abiraterone, especially in patients with cardiac comorbidities (15).

In the review of the literature, enzalutamide-induced non-ischemic cardiomyopathy was encountered only in two previously published case reports (8).

Case Presentation

We present a case of a 75-year-old Caucasian man who was diagnosed with non-

metastatic prostate cancer in 2010. He was experiencing dysuria for a couple of months before he scheduled an appointment with a urology specialist. The level of serum PSA was determined, and it was increased with an absolute value of 6.9 ng/ml. The urologist decided to perform a radical prostatectomy, as a primary diagnostic and treatment procedure, and it took place in December 2010. The pathology report indeed confirmed high-grade adenocarcinoma of the prostate gland, Gleason score 6, grade group 2, pT3NO with positive lymphovascular invasion, impaired capsule, and infiltration of adipose tissue as well as seminal vesicles. Due to tumor size and impaired capsule with infiltration of surrounding tissue, the tumor board decided to conduct postoperative radiation therapy. After completion of the treatment, in May 2011, with a total of 45 Gy of radiation dose, the patient started a watchful waiting regimen. In May 2019, at his regular 6-month follow-up, an increased PSA of 12.76 ng/ml was noted. Radiological findings, including thoracic, abdominal, and pelvic computed tomography scans and bone scintigraphy, were negative for loco-regional and distant metastases. He started with androgen deprivation therapy (LHRH agonist), and his PSA level decreased. Nevertheless, in March 2024, the PSA level started to rise again, with an absolute value of 44.29 ng/ml. Imaging studies showed enlarged retroperitoneal lymph nodes (up to 32 mm in diameter) and bone metastases. Since he had a medical history of well-controlled hypertension and atrial fibrillation, and had undergone surgery for an aneurysm of the abdominal aorta back in 1997, echocardiography was performed before making a therapy decision. Initial findings were within normal, with an ejection fraction of 61% so we have decided to initiate enzalutamide along with LHRH agonist treatment. The patient started enzalutamide therapy in July 2024.

In August 2024, 40 days after the initiation of enzalutamide therapy, he presented to the emergency room with an acute onset of shortness of breath, pain in the middle of the chest, peripheral edema, and fatigue. An echocardiogram demonstrated new cardiomyopathy with a reduced left ventricular (LV) ejection fraction of 26% and diffuse hypokinesis. Coronary angiography showed no significant arteriosclerosis. Enzalutamide was discontinued because it was the only preceding event before the onset of heart failure. He started guided heart failure therapy with beta-blocker (bisoprolol), angiotensin-converting enzyme inhibitor (ramipril), mineralocorticoid receptor antagonist (spironolactone), and sodium-glucose transport protein inhibitor (dapagliflozin). A control echocardiogram after one month showed improvement in left ventricular systolic function with an ejection fraction of 36%. Four months after enzalutamide discontinuation, the ejection fraction was 39%. The patient was still

experiencing fatigue, but less intense compared to the onset of the symptoms. The tumor board decided to continue LHRH agonist as monotherapy since the patient was not symptomatic from his metastatic cancer and due to the possible cardiotoxicity of other hormonal and chemotherapy agents.

Discussion

In the majority of cancer patients, death is caused by reasons other than malignancy itself. Around 45% of prostate cancer patients die due to complications of anticancer therapy (16). In recent decades, prostate cancer treatment has advanced significantly, with the approval of numerous new therapies, including hormonal treatments (17). Resistance to androgen deprivation therapy prompted the development of novel androgen receptor blockers, such as enzalutamide, apalutamide, and darolutamide, which have proven effective in the CRPC setting. These agents primarily work by inhibiting the interaction between androgens and AR, preventing AR nuclear translocation, or blocking AR-dependent gene transcription. These mechanisms ultimately reduce prostate cancer cell proliferation and tumor size but increase the possibility of the development of serious cardiovascular complications (18). Increased cardiovascular risk is reported in prostate cancer patients treated with ADT, abiraterone or enzalutamide. Hypermineralocorticism can be induced by abiraterone, which further leads to hypokalemia, hypertension, and edema. As a result of that, heart failure and atrial tachyarrhythmia may occur in patients treated with abiraterone (9).

Enzalutamide, an androgen receptor inhibitor, exhibits a higher binding affinity to androgen receptors compared to older inhibitors like bicalutamide. This characteristic is linked to improved clinical outcomes in patients with non-metastatic or metastatic castration-resistant prostate cancer compared to bicalutamide (19). Pathophysiology of QT prolongation, the most common cardiac adverse event of enzalutamide, is delayed rectifier potassium and sodium current, longer action potential duration, and appearance of afterdepolarizations (20). In a recent meta-analysis of prospective studies, enzalutamide was found to have a much better cardiac safety profile compared to abiraterone (21). A study done in 2021 showed a higher incidence of cardiac events in patients treated with abiraterone in contrast to the patients treated with enzalutamide and concluded that enzalutamide has a much better safety profile than abiraterone, especially in patients with cardiac comorbidities (15). Nevertheless, the patient did have cardiac adverse events due to enzalutamide therapy. In the review of the literature, enzalutamide-induced heart failure with reduced ejection fraction was encountered in only two previously reported cases.

A meta-analysis from 2018, which included 8,660 patients with metastatic prostate cancer treated with enzalutamide, found an increased incidence of hypertension among those patients (22). Another study by Shrestha B confirmed that the most common cardiovascular complication in patients treated with enzalutamide was hypertension (10.6%), followed by ischemic heart disease (1.88%) and atrial fibrillation (0.39%) (23).

Given the relatively high prevalence of cardiovascular complications associated with prostate cancer treatment, every patient must undergo a thorough clinical evaluation before initiating any anticancer therapy, including novel hormonal treatments. Treatment decisions should consider the patient's age, type of therapy, current and prior cardiovascular status, existing comorbidities, and concurrent medications. Regular cardio-oncology assessments are highly recommended, and existing cardiovascular conditions should be effectively managed and treated. Before starting therapy, patients should be screened for hypertension, dyslipidemia, and prediabetes/diabetes. Baseline electrocardiography and transthoracic echocardiography should be performed, with follow-up evaluations conducted before each new treatment phase. In the event of cardiovascular complications, management decisions should take into account the cancer prognosis (e.g., early-stage vs. metastatic disease), life expectancy, and the patient's preferences (24).

Conclusion

Prostate cancer has a very high prevalence among elderly men, and it could potentially be even higher due to the increasing elderly population worldwide. Anti-androgen therapy remains the standard of care in the majority of PC patients. In the past decade, several new hormonal agents were introduced in the treatment of prostate cancer. Although very efficient, new hormonal therapy is associated with an increased risk of many complications, of which cardiovascular complications are the most severe. One of the proposed mechanisms of the onset of hypertension, heart failure, ischemic heart disease, rhythm disturbances, and venous thromboembolic disease is the exacerbation of the imbalance of the cardiovascular system due to hormonal and metabolic changes. Hereby, we report a case of likely rare cardiovascular toxicity due to enzalutamide therapy, which typically has a safer cardiovascular profile among novel AR blockers. Literature review shows that abiraterone has more cardiovascular side effects than enzalutamide, which more commonly causes hypertension. Further research on the safety profile of novel anti-androgen therapy, as well as their impact on the cardiovascular system, is needed and should encourage the development of comprehensive cardio-oncology programs. Our

findings underline the importance of baseline screening and personalized evaluation of the cardiovascular system, risk stratification for possible cardiac adverse events, fundamental analysis of potential risk or benefit, before therapy initiation.

References

- Connolly RM, Carducci MA, Antonarakis ES. Use of androgen deprivation therapy in prostate cancer: indications and prevalence. *Asian J Androl* 2012; 14(2): 177-86. [\[CrossRef\]](#) [\[PubMed\]](#)
- Crawford ED, Heidenreich A, Lawrence N, Tombal B, Pompeo A, Mendoza-Valdes A, et al. Androgen-targeted therapy in men with prostate cancer: evolving practice and future considerations. *Prostate Cancer Prostatic Dis* 2019; 22(1): 24-38. [\[CrossRef\]](#) [\[PubMed\]](#)
- Živković-Perišić S, Miljuš D, Jovanović V. Registration of malignant diseases in estimating global cancer burden. *Glasnik javnog zdravljia* 2021; 95(1): 73-84.
- Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Cancer Stat Facts: Prostate Cancer, Aggregated With State, Total U.S. (2018-2023) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, released February 2025. Underlying mortality data provided by NCHS ([www.cdc.gov/nchs](#)).
- Mohler JL, Antonarakis ES. NCCN guidelines updates: management of prostate cancer. *J Natl Compr Canc Netw* 2019; 17(5.5): 583-6. [\[CrossRef\]](#) [\[PubMed\]](#)
- Fizazi K, Tran N, Fein L. Abiraterone acetate plus prednisone in patients with newly diagnosed high-risk metastatic castration-sensitive prostate cancer (LATITUDE): final overall survival analysis of a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2019; 20(5): 686-700. [\[CrossRef\]](#) [\[PubMed\]](#)
- de Bono JS, Logothetis CJ, Molina A, Fizazi K, North S, Chu L, et al. Abiraterone and increased survival in metastatic prostate cancer. *N Engl J Med* 2011; 364(21): 1995-2005. [\[CrossRef\]](#) [\[PubMed\]](#)
- Kumar A, Reddy A, Sekar A. Enzalutamide induced non-ischemic cardiomyopathy. A case report and review of literature on anti-androgen therapy-related cardiovascular events. *Cardiooncology* 2023; 9(1): 9. [\[CrossRef\]](#) [\[PubMed\]](#)
- Bretagne M, Lebrun-Vignes B, Pariente A, Shaffer CM, Malouf GG, Dureau P, Potey C, Funck-Brentano C, Roden DM, Moslehi JJ, Salem JE. Heart failure and atrial tachyarrhythmia on abiraterone: A pharmacovigilance study. *Arch Cardiovasc Dis* 2020; 113(1): 9-21. [\[CrossRef\]](#) [\[PubMed\]](#)
- Janssen. ZYTIGA (abiraterone acetate): highlights of prescribing information. Food and Drug Administration 2018; Available from: URL: https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/210308s000lbl.pdf
- Pfizer. XTANDI (enzalutamide): highlights of prescribing information. Food and Drug Administration 2019; Available from: URL: https://www.accessdata.fda.gov/drugsatfda_docs/label/2012/203415lbl.pdf
- Culig Z. Molecular mechanisms of enzalutamide resistance in prostate cancer. *Curr Mol Biol Rep* 2017; 3(4): 230-5. [\[CrossRef\]](#) [\[PubMed\]](#)
- Hussain M, Fizazi K, Saad F, Rathenborg P, Shore N, Ferreira U, et al.: Enzalutamide in Men with Nonmetastatic, Castration-Resistant Prostate Cancer. *N Engl J Med* 2018; 378(26): 2465-74. [\[CrossRef\]](#) [\[PubMed\]](#)
- Loriot Y, Miller K, Sternberg CN, Fizazi K, De Bono SJ, Chowdhury S, et al. Effect of enzalutamide on health-related quality of life, pain, and skeletal-related events in asymptomatic and minimally symptomatic, chemotherapy-naïve patients with metastatic castration-resistant prostate cancer (PREVAIL): results from a randomised, phase 3 trial. *Lancet Oncol* 2015; 16(5): 509-21. [\[CrossRef\]](#) [\[PubMed\]](#)
- Cone EB, Reese S, Marchese M, Nabi J, McKay RR, Kilbridge KL, Trinh QD. Cardiovascular toxicities associated with abiraterone compared to enzalutamide-A pharmacovigilance study. *EClinicalMedicine* 2021; 36: 100887. [\[CrossRef\]](#) [\[PubMed\]](#)
- Beyer DC, McKeough T, Thomas T. Impact of short course hormonal therapy on overall and cancer specific survival after permanent prostate brachytherapy. *Int J Radiat Oncol Biol Phys* 2005; 61(5): 1299-1305. [\[CrossRef\]](#) [\[PubMed\]](#)
- Wilk M, Waśko-Grabowska A, Szmit S. Cardiovascular Complications of Prostate Cancer Treatment. *Front Pharmacol* 2020; 11: 555475. [\[CrossRef\]](#) [\[PubMed\]](#)
- Rice MA, Malhotra SV, Stoyanova T. Second-generation antiandrogens: from discovery to standard of care in castration resistant prostate cancer. *Front Oncol* 2019; 9: 801. [\[CrossRef\]](#) [\[PubMed\]](#)
- Penson DF, Armstrong AJ, Concepcion R, Agarwal N, Olsson C, Karsh L, et al. Enzalutamide versus bicalutamide in castration-resistant prostate cancer: the STRIVE Trial. *J Clin Oncol* 2016; 34(18): 2098-106. [\[CrossRef\]](#) [\[PubMed\]](#)

20.Saigal CS, Gore JL, Krupski TL, Hanley J, Schonlau M, Litwin M, et al. Androgen deprivation therapy increases cardiovascular morbidity in men with prostate cancer. *Cancer* 2007; 110(7): 1493–500. [\[CrossRef\]](#) [\[PubMed\]](#)

21.Salem JE, Yang T, Moslehi JJ, Waintraub X, Gandjbakhch E, Bachelot A, et al. Androgenic effects on ventricular repolarization: a translational study from the international pharmacovigilance database to iPSC-cardiomyocytes. *Circulation* 2019; 140(13): 1070–80. [\[CrossRef\]](#) [\[PubMed\]](#)

22.Lacovelli R, Ciccarese C, Bria E, Romano M, Fantinelli E, Bimbatti D, et al. The cardiovascular toxicity of Abiraterone and Enzalutamide in prostate cancer. *Clin Genitourin Cancer* 2018; 16(3):e645–53. [\[CrossRef\]](#) [\[PubMed\]](#)

23.Shrestha B, Gouli S, Shrestha A. Abstract 16180: Cardiovascular adverse events associated with androgen receptor-targeted therapy used in the treatment of prostate cancer (abstract). *Circulation* 2020; 142:A16180. [\[CrossRef\]](#) [\[PubMed\]](#)

24.Curigliano G, Lenihan D, Fradley M, Ganatra S, Barac A, Blaes A, et al. Management of cardiac disease in cancer patients throughout oncological treatment: ESMO consensus recommendations. *Ann Oncol* 2020; 31(2): 171–90. [\[CrossRef\]](#) [\[PubMed\]](#)

Prikaz bolesnika

UDC: 616.12-008.46:615.277

doi: 10.5633/amm.2025.0315

SRČANA INSUFICIJENCIJA SA REDUKOVANOM EJEKCIJONOM FRAKCIJOM UZROKOVANA ENZALUTAMIDOM

*Mirjana Todorović Mitić¹, Marija Todorović Marković², Ana Cvetanović^{1,3},
Milan Čevrljaković¹, Ljubomir Dinić^{4,5}*

¹Univerzitetski klinički centar Niš, Klinika za onkologiju, Niš, Srbija

²Univerzitalska bolnica Rigshospitalet (Visokospecijalizovana bolnica u Danskoj), Klinika za kardiologiju, Kopenhagen, Danska

³Univerzitet u Nišu, Medicinski fakultet, Katedra Onkologija, Niš, Srbija

⁴Univerzitet u Nišu, Medicinski fakultet, Katedra Hirurgija i anestezijologija sa reanimatologijom, Niš, Srbija

⁵Univerzitetski klinički centar Niš, Klinika za urologiju, Niš, Srbija

Kontakt: Mirjana Todorović Mitić
Zorana Radosavljevića Čupe 2a, 18000 Niš, Srbija
E-mail: mirjanatdrvc@gmail.com

Karcinom prostate je drugo najčešće maligno oboljenje kod muškaraca i drugi vodeći uzrok smrti od malignih bolesti. Lečenje inoperabilnih pacijenata, pacijenata sa lokalno uznapredovalom ili metastatskom bolesti zahteva primenu terapije derivacije androgena.

U protekloj deceniji u terapiju karcinoma prostate uvedeni su lekovi enzalutamid i abirateron. Bezbednosni profil ovih lekova i podaci iz kliničke prakse ukazuju na to da pomenuti lekovi imaju značajan uticaj na pojavu kardiovaskularnih komplikacija. Srčana insuficijencija je češći neželjeni efekat primene abiraterona. U ovom radu se prikazuje slučaj pacijenta kod kojeg je pojavu srčane insuficijencije izazvao enzalutamid. Pregledom literature nađena su samo dva slučaja srčane insuficijencije uzrokovane enzalutamidom u svetu.

Neophodna su dalja istraživanja o uticaju nove antiandrogene terapije na kardiovaskularni sistem, kao i personalizovani pristup proceni funkcije kardiovaskularnog sistema i stratifikacija rizika od pojave potencijalnih kardiovaskularnih neželjenih efekata pre započinjanja terapije.

Acta Medica Mediana 2025; 64(3): 118–123.

Ključne reči: enzalutamid, srčana insuficijencija, karcinom prostate

"This work is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) Licence".