

**CHEMICAL COMPOSITION AND ANTIOXIDANT ACTIVITY OF NOBLE YARROW
(*ACHILLEA NOBILIS* L. SUBSP. *NEILREICHII* (KERNER) VELEN.)**

**Katarina Radovanović¹, Maja Hitl¹, Biljana Božin^{1,2}, Katarina Bijelić¹,
Blagoje Prpa¹, Milica Aćimović³, Neda Gavarić^{1,2*}**

¹University of Novi Sad – Faculty of Medicine, Department of Pharmacy, Novi Sad,
Serbia

²Center for Medical and Pharmaceutical Investigations and Quality Control, Novi
Sad, Serbia

³Institute of Field and Vegetable Crops, Novi Sad, Serbia

*neda.gavaric@mf.uns.ac.rs

Species of the genus *Achillea* are well-known ancient medicinal plants. Noble yarrow is presented in eastern Serbia, Suva planina and Kosovo (around Peja). Due to the traditional use, it is important to determine the chemical composition and biological activities of these species widespread in Serbia. The aim of this study was to determine the chemical composition and antioxidant activity of noble yarrow. The content of total phenols in the prepared infusion was determined by the Folin – Ciocalteu method. Total flavonoids were determined spectrophotometrically by forming complexes with AlCl₃. Chemical characterization was performed by high performance liquid chromatography. The antioxidant activity was examined spectrophotometrically by determining the "scavenger" activity of 2,2-diphenyl-1-picrylhydrazyl (DPPH), hydroxyl (OH •) radical and nitroso (NO •) radical, as well as redox potential testing by FRAP test and determination of inhibition lipid peroxidation in the Fe²⁺ / H₂O₂ induction system. The total phenol content in the tested infusion was 33.52 mg of gallic acid equivalent per gram of dry extract, while the content of total flavonoids was 6.79 mg of quercetin equivalent per gram of dry extract. Phenolic acids are dominantly presented in the infusion. Rosmarinic and *p*-coumaric acids are the most abundant (13.34 and 12.74 mg/g of dry extract). The infusion showed the ability to neutralize free radicals at medium high concentrations. Unlike the results of other antioxidant tests, inhibition of lipid peroxidation was not achieved. The tested plant is a good source of phenolic compounds with established antioxidant activity.

Acknowledgements

The Ministry of Education, Science and Technological Development, Republic of Serbia, Grant Number 451-03-68/2022-14/200114

HEMIJSKI SASTAV I ANTIOKSIDANTNA AKTIVNOST ŽUĆKASTOG SPORIŠA (*ACHILLEA NOBILIS* L. SUBSP. *NEILREICHII* (KERNER) VELEN.)

**Katarina Radovanović¹, Maja Hitl¹, Biljana Božin^{1,2}, Katarina Bijelić¹,
Blagoje Prpa¹, Milica Aćimović³, Neda Gavaric^{1,2*}**

¹Univerzitet u Novom Sadu – Medicinski fakultet, Katedra za farmaciju, Novi Sad,
Srbija

² Centar za medicinsko-farmaceutska istraživanja i kontrolu kvaliteta, Novi Sad,
Srbija

³Institut za ratarstvo i povrtarstvo, Novi Sad, Srbija

* neda.gavaric@mf.uns.ac.rs

Vrste roda *Achillea* su široko rasprostranjene lekovite biljke sa poznatom tradicionalnom primenom od davnina. *Achillea nobilis* L. subsp. *neilreichii* (Kerner) Velen., u narodu poznata kao žućkasti sporiš, rasprostranjena je na teritoriji istočne Srbije, Suve planine i Kosova (okolina Peći). Zbog raširene tradicionalne primene, značajno je da se detaljno prouče hemijski sastav i biološke aktivnosti vrsta roda *Achillea* rasprostranjenih u Srbiji, pa je cilj ovog rada bio da se odrede hemijski sastav i antioksidantna aktivnost infuza žućkastog sporiša. U pripremljenom infuzu određen je sadržaj ukupnih fenola metodom po *Folin-Ciocalteu*. Ukupni flavonoidi određeni su spektrofotometrijski praćenjem formiranja kompleksa sa AlCl₃. Hemiska karakterizacija izvršena je visokoefikasnom tečnom hromatografijom. Antioksidativno delovanje infuza ispitano je spektrofotometrijski određivanjem „skevindžer“ aktivnosti na 2,2-difenil-1-pikrilhidrazil (DPPH), hidroksil (OH•) radikal i nitrozo (NO•) radikal, kao i ispitivanjem redoks potencijala pomoću FRAP testa i određivanjem inhibicije lipidne peroksidacije u Fe²⁺/H₂O₂ sistemu indukcije. Sadržaj ukupnih fenola u ispitivanom infuzu je 33,52 mg ekvivalenta galne kiseline po gramu suvog ekstrakta, dok je sadržaj ukupnih flavonoida 6,79 mg ekvivalenta kvercetina po gramu suvog ekstrakta. U infuzu su dominantno prisutne fenolne kiseline od kojih su najzastupljenije rozmarinska (13,34 mg/g suvog ekstrakta) i *p*-kumarna kiselina (12,74 mg/g suvog ekstrakta). Infuz je pokazao sposobnost da neutralizuje slobodne radikale pri srednje visokim koncentracijama. Za razliku od rezulata ostalih antioksidativnih testova, pomoću analiziranog infuza nije postignuta inhibicija lipidne peroksidacije. Ispitivana biljka predstavlja dobar izvor fenolnih jedinjenja sa utvrđenim antioksidantnim delovanjem.

Zahvalnica

Ministarstvo prosvete, nauke i tehnološkog razvoja Republike Srbije, Projekat: 451-03-68/2022-14/200114