

## A STUDY ON NUTRITIONAL STATUS AMONG THE ADULT GAROS' OF WEST GARO HILLS, NORTH-EAST INDIA (MEGHALAYA)

**Khriesivonuo Nakhro, Kusum Ghosh, Diptendu Chatterjee, Arup R Bandyopadhyay**

Department of Anthropology, University of Calcutta, Kolkata-700019, West Bengal, India

**Abstract.** Health Status is the basic indicator of human wellbeing and nutritional status. Deficiencies or imbalance in nutrient intake leads to malnutrition which is associated with a number of short- and long-term health consequences that alleviate the ability of an individual to sustain economically productive work and ultimately hinder the development of the whole nation. Thus, study on nutritional status requires urgent attention in order to prevent malnutrition related complications. In this background, the present study examines nutritional status of the Garos of West Garo Hills, Meghalaya. To achieve this purpose, cross-sectional study was conducted among 43 females and 57 males of 'Garos' from Asananggre village, West Garo Hills, North-east India (Meghalaya). The study divulge taller and heavier males compared to females. Body Mass Index status of male and female did not differ significantly ( $p < 0.05$ ). Majority of females belong to normal (52.6%) followed by overweight category (35.1%) while males were more incline to overweight category (48.8%). According to MUAC, the nutritional status of both the males and females are normal. Significant ( $p < 0.05$ ) difference was seen between males and females in terms of Hemoglobin level where, majority were mild anemic or normal. 1.8% females were found to be severely anemic. From the study it can be envisaged that most adult males and females are well-nourished and only few females were severely anemic i.e., undernourished.

**Key words:** health, malnutrition, hemoglobin level, BMI, MUAC, anemia

### Introduction

Health status is central to human happiness, well-being and resource quality (Odonkor et al., 2019) while nutritional status is an important determinant for maintaining optimum health (Jumrakh et al., 2001; Muslim et al., 2021) being the prime indicator of health status. Human body require proper nutrition through well balanced diet to fulfill body needs and to maintain basic physiological functions of the body. Therefore any deficiencies or imbalance in nutrient intake leads to malnutrition which can be either 'overnutrition', consumption of excess calorie or 'undernutrition', insufficient supply of one or more essential nutrients (Bhattacharya et al., 2019).

Both undernutrition and overnutrition is consequently the most important risk factor for the burden of disease throughout the world, particularly in developing countries (Paruchuri et al., 2012; Muller et al., 2005; Nemer et al., 2001), affecting 792 million people worldwide, of which 20% are from developing countries (World Bank Report, 2011). Over-nutrition is a threat that increases body weight and causes several non-communicable diseases such as, metabolic syndrome, high cholesterol, type 2 diabetes mellitus, high blood pressure, and cardiovascular disease which are a threat to the public health in the developing countries like India (Little et al., 2016; Bhattacharya et al., 2019; Muslim et al., 2021). On the other hand, it is well documented that, undernutrition results in both short- and long-term health effects

---

Autor za korespondenciju: Khriesivonuo Nakhro, e-mail: khriesivonuoakhro@gmail.com  
Department of Anthropology, University of Calcutta, 35 Ballygunge Circular Road, Kolkata-700019, West Bengal, India

Primljeno u redakciju 17. 8. 2021, revidirana verzija 22. 8. 2021, rad prihvaćen za publikovanje 19. 9. 2011.

and even death by decreasing body immunity (Maleta, 2006; Kuruvilla et al., 2016). The short-term effect of undernutrition includes, weakness and recurring illness, which ultimately, in the long run hampers all vital physiological functions of the body causing measurable effect on body composition i.e., low weight, growth retardation in children (stunting, wasting, Kwashiorkor and Marasmus) and adolescent. Furthermore, undernutrition induced decreased immunity leading to recurring infections (Muller et al., 2005) and also the occurrence of chronic diseases like diabetes mellitus, hypertension, coronary heart diseases (Hoet, 1997), anemia (Thakur et al., 2014) and impaired mental development (Martins et al., 2011) in later adult life. Moreover, among women, undernutrition may cause obstetric complications leading to maternal and infant mortality and also increases the probability of giving low birth weight babies (Maleta, 2006; Rotimi et al., 1999; Allen et al., 1994) and thus starts the undernutrition cycle again, spanning several generations. Beside this, chronic undernutrition also alleviates the ability to sustain economically productive work by reducing working capacity of an individual which ultimately results in low income (Roy, 2002; Stewart et al., 2003). Therefore, undernutrition is not only a health burden but an economic burden affecting development of human beings as well as the whole nation.

Anthropometry being one of the methods used to assess the nutritional status, is widely used for its non-invasive, inexpensive and provides detailed information on different components of body structure especially muscular and fat components (Bharati et al., 2007; Bhattacharya et al., 2019). Studies reported (Szczepanska et al., 2019; Bhattacharya et al., 2019), among the widely used anthropometric measurements, body mass index (BMI) and mid-upper-arm-circumference (MUAC) as most significant and reliable one. Besides anthropometric measurements, hemoglobin (Hb) level is another important indicator of improper nutritional condition (McLean et al., 2007; Mondini et al., 2009) as it is associated with both macro- and micronutrient deficiencies including iron, vitamin B12 and folic acid deficiency, considered as the major cause of low hemoglobin level i.e., anemia (Emiroglu et al., 2019).

The Northeastern region of India is inhabited by a number of small and large tribes. Meghalaya is one of the eight sisters states in the Northeastern part of India, comprising Khasi, Garo and Jaintia as the three main tribes of Meghalaya (Singh et al., 2019). Due to poor healthcare coverage and lack of healthcare infrastructure, all the northeastern states have been given particular focus under the National Rural Health Mission (NRHM) (Meitei, 2020) and Prime Minister's 20-point program (Gopalan, 1992). A trend of developmental transition is underway in India and its Northeastern parts (Amuna et al., 2008, Sikdar, 2008; Sikdar, 2012), in all aspects of human life including disease pattern, demographic pattern, nutritional status, socioeconomic and ecological conditions (Dasgupta et al., 2008) but the progress is uneven and inequitable. Therefore, it is necessary to design region-specific public health policies (Yu et al., 2019) to find early intervention strategies and mitigate the human resource and economic burden of those regions and ultimately to India by investigating the nutritional status. Several studies are there in this regard in different regions (Swain et al., 2018; Bhattacharya et al., 2019) on different age groups (Kanjilal et al., 2010; Agarwalla et al., 2015; Padma et al., 2016) but most of the previous studies on nutritional status from Meghalaya, North-east India and Garos of East Garo Hills are conducted on children (Rao et al., 2005; Chyne et al., 2017; Singh et al., 2019; Meitei, 2020) and a very few on women (Chyne et al., 2017; Nongrum et al., 2021) while, research related to the prevalence of malnutrition among the adults (men and women) of North-eastern part of India, particularly from the Garo tribes of Meghalaya are meagre.

## Objectives

So, in this backdrop the present study has been undertaken, to investigate the nutritional status of the adult Garos, living in Asananggre Village of West Garo Hills, Meghalaya (North-east India).

## Materials and Methods

The present study comprises of 100 adult individuals including both male and female (non-pregnant), from 152 households of Asananggre Village, West Garo Hills, Meghalaya (Figure 1). Prior to study verbal consent were obtained from the participants. Data has been collected on anthropometric variables (Height and Weight) by using standard technique (Weiner and Lourie, 1981). Then by using these anthropometric measures required indices (BMI) were calculated and classified according to WHO classification of BMI for Asian adults 2013. Mid-calf circumference and mid-upper arm circumference was taken by putting the tape around the most developed muscles and categorized using WHO classification of MUAC 1995.



**Figure 1** Map of West Garo Hills in Meghalaya, North-east India

Hemoglobin concentration was estimated by Sahli's method. The tube is filled with 2 grams marked with N/10 HCl. Blood was collected using finger prick method, using a disposable needles and blood was drawn into Sahli's pipette up to 20 micro liter marking. HCl and blood were stirred, the solution is left for 5 minutes to form acid haematin. Then the acid haematin is diluted by adding distilled water drop by drop till it matched the standard color plates of the comparator. Results were read as gram/dl and classified according to WHO, 1975. Necessary descriptive and inferential statistics are calculated by using SPSS version 18.

## Results

Present study envisaged that, the males are taller and heavier than the females (Table 1). Not only that, the nutritional markers also vindicated higher values in case of the males. Furthermore, Hemoglobin level is lower in females than males. According to BMI (Table 2) maximum number of females belong to normal category (52.6%) followed by overweight (35.1%) and underweight (12.3%). But males were prone to overweight (48.8%). BMI status did not differ significantly ( $p < 0.05$ ) between males and females. Distribution of MUAC (Table 3) envisaged that, the nutritional status of both the males and females were normal.

Significant ( $p < 0.05$ ) differences were found between males and females in terms of Hemoglobin level (Table 4). Most of the males and females are mild anemic or normal and only 1.8% females belong to severe anemic category.

**Table 1.** Distribution of anthropometric variables among the males and females

Variables	Males (43) (Mean $\pm$ SD)	Females (57) (Mean $\pm$ SD)
Height	162.6 $\pm$ 7.4	149.4 $\pm$ 5.4
Weight	60.6 $\pm$ 10.9	51.9 $\pm$ 10.1
Mid-upper arm Circumference	26.2 $\pm$ 2.7	24.9 $\pm$ 2.9
Mid-calf Circumference	35.3 $\pm$ 2.8	33.5 $\pm$ 3.5
Body Mass Index	22.7 $\pm$ 2.9	22.3 $\pm$ 3.1
Hemoglobin Level	13.2 $\pm$ 1.6	11.1 $\pm$ 1.7

The results of the linear regression analysis (Table 2) indicate that together all examined socioeconomic and demographic factors significantly affect BMI values and together explain 9% of variability. Gender, age, mother's education and the number of children in the family have the most significant influence on BMI.

**Table 2.** Distribution of Body Mass Index (BMI) among adult males and females

Category	Underweight (%)	Normal (%)	Overweight	Total Frequency (100%)	Chi-square value
Males	2 (4.7)	20 (46.5)	21 (48.8)	43	0.87
Females	7 (12.3)	30 (52.6)	20 (35.1)	57	

**Table 3.** Distribution of Mid Upper Arm Circumference (MUAC) among adult males and females

Category	Male (%)	Category	Female (%)
Undernourished (<23)	4 (9.3)	Undernourished (<22)	7 (12.3)
Normal ( $\geq$ 24)	39 (90.7)	Normal ( $\geq$ 23)	50 (87.7)

**Table 4.** Distribution of hemoglobin level among adult males and females

Category	Severe Anemia (%)	Moderate Anemia (%)	Mild Anemia (%)	Normal (%)	Total Frequency (100%)	Chi-square value
Males	0	0	16 (37.2)	27 (62.8)	43	16.1*
Females	1 (1.8)	16 (28.0)	18 (31.6)	22 (38.6)	57	

\*significant  $p < 0.05$

## Discussion

Malnutrition is associated with both short (weakness and recurring illness) and long-term health consequences (low weight, growth retardation, infections, low birth weight baby and non-communicable diseases) and even death by decreasing body immunity (Muller et al., 2005; Bhattacharya et al., 2019; Muslim et al., 2021; Hoet, 1997; Thakur et al., 2014; Maleta, 2006; Kuruvilla et al., 2016) which can pass on to several generations. Undernutrition is not only a health burden but an economic burden too, (Roy, 2002; Stewart et al., 2003) that hinder the development of the whole nation. Thus, study on nutritional status requires urgent attention in order to prevent malnutrition related health outcomes.

The present study envisaged maximum number of the females belonging to normal weight category followed by overweight. These corroborated with earlier studies done among two ethnic groups of Tripura, Northeast India (Bhattacharya et al., 2019; Ghosh et al., 2018) and Karbi Women of Assam, Northeast India (Little et al., 2016). This may be due to nutritional transition because of an interplay among economic, demographic, environmental and cultural changes occurring in the society (Swain et al., 2018). But in case of males, they were found to be more prone to the overweight category, which was in contrast with the previous studies (Rengma et al., 2015; Mungreiphy et al., 2008) done in the regions of North-east India, which reported higher prevalence of overweight in female than males. Furthermore, the present study also comprehensively indicates that, nutritional status of most of the males and females are normal in terms of both MUAC and hemoglobin level, only 1.8% females belong to severe anemic category. Looking into the hemoglobin level, the present study reveal females having more tendency to be anemic compared to male. Similar findings also reported in the previous studies (Azad et al., 2017, Bentley et al., 2003) done in India and in districts of Meghalaya (Mongsang et al., 2018).

Therefore, from the foregoing study we can conclude that, most of the males and females of Asananggre Village, West Garo Hills, Meghalaya are well-nourished and only few of the females are severely anemic i.e., undernourished. Therefore, further prospective studies are needed on the factors that instigate the low level of hemoglobin among females, which could help in taking appropriate measures for preclusion and in promoting the health status of the individuals of the studied area. The present study attempted, also implies that more such studies are needed in other ethnic groups of North-eastern part of India to find out who are at the higher risk and need an immediate intervention Program.

**Acknowledgement.** This financially independent study would not have been possible without the participants, therefore sincere thanks to all the participants for their contribution and co-operation. Authors further extend acknowledgement to Dr. Sudipta Ghosh for her immense help and guidance.

## References

- Agarwalla, R., Saikia, A.M., Baruah, R. (2015). Assessment of the nutritional status of the elderly and its correlates. *Journal of Family and Community Medicine*, 22(1), 39–43.
- Allen, L.H., Lungaho, M.S., Shaheen, M., Harrison, G.G., Neumann, C., Kirksey, A. (1994). Maternal body mass index and pregnancy outcome in the nutrition collaborative research support program. *European Journal of Clinical Nutrition*, 48, S68–76.
- Amuna, P., Zotor, F.B. (2008). Epidemiological and nutrition transition in developing countries: Impact on human health and development. *Proceedings of Nutrition Society*, 67, 82–90.
- Azad, K.L., Dwivedi, S.K. (2017). A study of profile of patients with different degrees of anaemia. *International Journal of Research in Medical Sciences*, 5(9), 4008–4012.
- Bentley, M.E., Griffiths, P.L. (2003). The burden of anaemia among women in India. *European Journal of Clinical Nutrition*, 57, 52–60.
- Bharati, S., Pal, M., Bhattacharya, B.N., Bharati, P. (2007). Prevalence and causes of chronic energy deficiency and obesity in Indian women. *Human Biology*, 79, 395–412.
- Bhattacharya, A., Pal, B., Mukherjee, S., Roy, S.K. (2019). Assessment of nutritional status using anthropometric variables by multivariate analysis. *BMC Public Health*, 19(1045), 1–9.
- Bhattacharya, P., Chatterjee, D., Sarkar, P., Ghosh, K., Bandyopadhyay, A.R. (2019). Association of fat patterning, type 2 diabetes mellitus and MTHFR gene polymorphism: a study among the two ethnic groups of Tripura, North-East India. *European Journal of Clinical and Experimental Medicine*, 17(3), 209–213.

- Chyne, D.A.L., Meshram, I.I., Rajendran, A., Kodali, V., Getti, N., Roy, P., Kuhnlein, H.V., Longvah, T. (2017). Nutritional status, food insecurity, and biodiversity among the Khasi in Meghalaya, North-East India. *Maternal and Child Nutrition*, 13(3), e12557.
- Dasgupta, P., Saha, R., Nube, M. (2008). Changes in body size, shape and nutritional status of middle-class Bengali boys of Kolkata, India, 1982-2002. *Economics and Human Biology*, 6, 75-94.
- Emiroglu, C., Gorpelioglu, S., Aypak, C. (2019). The Relationship between Nutritional Status, Anemia and other Vitamin Deficiencies in the elderly Perceiving home care. *The Journal of Health, Nutrition and Aging*, 23(7), 677-682.
- Ghose, B., Tang, S., Yaya, S., Feng, Z. (2016). Association between food insecurity and anaemia among women of reproductive age. *Peer J*, 4, e1945, 1-12.
- Ghosh, K., Sarkar, P., Chatterjee, D., Bandyopadhyay, A.R. (2018). Association of fat patterning, hypertension and ACE (I/D) gene polymorphism: a study on two Tibeto Burman linguistic group of Tripura, North East India. *MOJ Anatomy & Physiology*, 5(6), 368-371.
- Gopalan, C. (1992). Nutrition challenge for Asia. *Progress in Food and Nutrition Science*, 16(1), 51-84.
- Hoet, J.J. (1997). The role of fetal and infant growth and nutrition in the causality of diabetes and cardiovascular disease in later life. *SCN News*, 14, 10-3.
- Jumrakh, M., Lubis, I.Z., Aziz, N. (2001). Nutritional status and hemoglobin level in elementary schoolchildren. *Paediatrica Indonesiana*, 41, 296-298.
- Kanjilal, B., Mazumdar, P.G., Mukherjee, M., Rahman, M.H. (2010). Nutritional status of children in India: household socio-economic condition as the contextual determinant. *International Journal for Equity in Health*, 9(19), 1-13.
- Kuruvilla, S., Bustreo, F., Kuo, T., Mishra, C.K., Taylor, K., Fogstad, H., Gupta, G.R., Gilmore, K., Temmerman, M., Thomas, J., Rasanathan, K., Chaiban, T., Mohan, A., Gruending, A., Schweitzer, J., Dini, H.S., Borrazzo, J., Fassil, H., Gronseth, L., Khosla, R., Cheeseman, R., Gorna, R., McDougall, L., Toure, K., Rogers, K., Dodson, K., Sharma, A., Seoane, M., Costello, A. (2016). The global strategy for women's, children's and adolescents' health (2016-2030): a roadmap based on evidence and country experience. *Bull World Health Organ*, 94, 398-400.
- Little, M., Humphries, S., Patel, K., Dewey, C. (2016). Socio-Economic and Demographic Correlates of Overweight and Obesity: A Study on the Karbi Women of Assam, Northeast India. *Anthropology Open Journal*, 2(2), 31-39.
- Maleta, K. (2006). Undernutrition. *Malawi Medical Journal*, 18, 189-205.
- Martins, V.J.B., Florencio, T.M.M.T., Grillo, L.P., Carmo, Franco, M.C.P., Martins, P.A., Clemente, A.P.G., Santos, C.D.L., Vieira, M.F.A., Sawaya, A.L. (2011). Long-lasting effects of undernutrition. *International Journal of Environmental Research and Public Health*, 8, 1817-46.
- McLean, E., Egli, I., Cogswell, M., Benoist, B., Wojdyla, D. (2007). Worldwide prevalence of anemia in preschool age children and pregnant women and non-pregnant women of reproductive age. In *Te guide book : Nutritional Anemia*. Basel, J.B., Zimmermann, M.B., & Kraemer, K. Eds., pp. 11-12. Switzerland: Sight and Life Press.
- Meitei, W.B. (2020). Nutritional Status of Under-five Children in Northeast India: Effect of Household Socioeconomic Status. *Indian Journal of Public Health Research & Development*, 11(7), 1003-1008.
- Mondini, L., Rodrigues, D.A., Gimeno, S.G.A., Baruzzi, R.G. (2009). Anemia and hemoglobin levels among Indigenous Xavante children, Central Brazil. *Rev Bras Epidemiol*, 12(3), 1-8.
- Monsang, E.S., Singh, N. (2018). Prevalence of Anaemia and Nutritional Knowledge among Tribal Women of Reproductive Age Group of Meghalaya, India. *International Journal of Current Microbiology and Applied Sciences*, 7(10), 1221-1229.
- Muller, O., Krawinkel, M. (2005). Malnutrition and health in developing countries. *Canadian Medical Association Journal*, 173(3), 279-286.
- Mungreiphy, N.K., Kapoor, S. (2008). Overweight, Obesity and Socio- economic change among Tankhul Naga tribal women of Manipur, North East India. *Nature Proceedings*, doi.org/10.1038/npre.2008.1947.1
- Muslim, A., Lim, Y.A., Sofian, S.M., Shaari, S.A., Zain, Z.M. (2021). Nutritional status, hemoglobin level and their associations with soil transmitted helminth infections between Negritos (indigenous) from the inland jungle village and resettlement at town peripheries. *Plos one* 16(1), e0245377

- Nemer, L., Gelband, H., Jha, P., & Commission on Macroeconomics and Health. (2001). *The evidence base for interventions to reduce malnutrition in children under five and school-age children in low- and middle-income countries*. CMH working paper no WG5:11. Geneva: World Health Organization.
- Nongrum, M.S., Pawera, L., Mawroh, B. (2021). Dietary diversity and its determinants among Khasi and Garo indigenous women (15 to 49 years) in Meghalaya, northeast India. *Nutrition and Health*, doi: 10.1177/02601060211016629
- Nowak-Szczepanska, N., Gomula, A., Koziel, S. (2019). Mid-upper arm circumference and body mass index as different screening tools of nutritional and weight status in Polish schoolchildren across socio-political changes. *Scientific reports*, 9, 12399.
- Odonkor, S.T., Kitcher, J., Okyere, M., Mahami, T. (2019). Self assessment of Hygiene practices towards predictive medicine intervention: A case study of University students in Ghana. *BioMed Research International*, doi:10.155/2019/3868537
- Padma, A.S., Sukumaran, M.K., Padma, S., Rajani, D., Vanitha, S. (2016). Evaluation of Nutritional Status of College Girl Students in terms of Haemoglobin levels and BMI. *Journal of Medical Science and Clinical Research*, 4(9), 12822-12827.
- Paruchuri, A., Vini, P., Ahmad, A., Manna, P.K., Kumaran, S., Mohanta, G.P. (2012). Malnutrition in India: A Major Problem with Minor Attention. *IJPI's Journal of Hospital and Clinical Pharmacy*, 2(9), 27-35.
- Rao, G.R., Ladusingh, L., Pritamjit, R. (2005). Nutritional status of children in north-east India. *Asia-Pacific Population Journal*, 19(3), 39 – 56.
- Rengma, M.S., Sen, J., Mondal, N. (2015). Socio – Economic, demographic and lifestyle determinants of overweight and obesity among adults of Northeast India. *Ethiopian Journal of Health Science*, 25(3), 199-208.
- Rotimi, C., Okosun, I., Johnson, L., Owoaje, E., Lawoyin, T., Asuzu, M., Kaufman, J., Adeyemo, A., Cooper, R. (1999). The distribution and mortality impact of chronic energy deficiency among adult Nigerian men and women. *European Journal of Clinical Nutrition*, 53, 734–9.
- Roy, P.P., Gunturu, V.V. (2017). Association of anaemia with BMI in Medical students: a cross-sectional study. *Journal of Dental and Medical Sciences*, 16(8), 32-35.
- Roy, S.K. (2002). Factors affecting the work productivity of Oraon agricultural laborers of Jalpaiguri district, West Bengal. *American Journal of Physical Anthropology*, 117, 228–35.
- Sikdar, M. (2008). Influence of socioeconomic transition on genetic structure: A case study in Upper Assam, India. *Annals of Human Biology*, 35, 112–20.
- Sikdar, M. (2012). Socioeconomic covariates and their impact on the opportunity for natural selection in a riparian tribe of Northeast India. *Anthropologischer Anzeiger*, 69, 273–87.
- Singh, N., Nagar, S., Takhellambam, R.D. (2019). Nutritional Status of Tribal Primary School Children in Meghalaya. *International Journal Current Microbiology and Applied Sciences*, 8(7), 668-681.
- Stewart, W.F., Ricci, J.A., Chee, E., Morganstein, D. (2003). Lost productive work time costs from health conditions in the United States: Results from the American productivity audit Health and Lost Productive Time in the US Workforce. *Journal of Occupational and Environmental Medicine*, 45, 1234–46.
- Swain, S., Chowdhury, S. (2018). Trends of nutritional status among rural adults in six states of India: findings from national survey data. *Clinical Epidemiology and Global Health*, 6, 181–187.
- Thakur, N., Chandra, J., Pemde, H., Singh, V. (2014). Anemia in severe acute malnutrition. *Nutrition*. 30(4), 440-2.
- Weiner, J.S., Lourie, J.A., (1981). *Practical Human Biology*. New York: Academic Press.
- World Bank Report on Malnutrition in India; World Bank Report. 2011. [cited 2021 june]. Available in: <http://www.worldbank.org/WBSITE/COUNTRIES/SOUTH/ASIAEXT/>
- Yu, S., Xing, L., Du, Z., Tian, Y., Jing, L., Yan, H., Lin, M., Zhang, B., Liu, S., Pan, Y., Li, C. (2019). Prevalence of Obesity and Associated Risk Factors and Cardio metabolic Comorbidities in Rural Northeast China. *BioMed Research International*, doi:10.1155?2019?6509083