

**LINC00467 INDUCES MELANOMA DETERIORATION BY
TARGETING MIR-485-5P/P21 ACTIVATED KINASE 1**LINC00467 IZAZIVA POGORŠANJE MELANOMA CILJANJEM MIR-485-5P/P21
AKTIVIRANJEM KINAZE 1Zhoujing Ji^{1#}, Jie Zhang^{1#}, Lili Zhang¹, Shengju Yang¹, Yangcheng Li^{2*}, Lixiong Gu^{1*}¹Department of Dermatology, Affiliated Hospital of Nantong University, Nantong, China²Department of General Surgery, Tumor Hospital Affiliated to Nantong University, Nantong Tumor Hospital,
Nantong, China**Summary****Background:** The purpose of the current research was to investigate the biological roles of LINC00467 in inducing melanoma deterioration.**Methods:** Differential level of LINC00467 in melanoma tissues and its prognostic value were analyzed in GEPIA, which were further confirmed in clinical samples we collected. Regulatory effects of LINC00467 on proliferation, migration and invasion capacities of A375 and SKMEL1 cell lines were examined by a series of functional experiments. Potential downstream targets of LINC00467 were identified through dual-luciferase reporter assay, and their synergistic role in melanoma process was finally explored by rescue experiments.**Results:** LINC00467 was up-regulated in melanoma samples, but it did not have a prognostic potential in melanoma. LINC00467 has the capacities to stimulate proliferation, migration and invasion of A375 and SKMEL1 cell lines. The feedback loop LINC00467/miR-485-5p/PAK1 was identified, which was responsible for inducing melanoma deterioration.**Conclusions:** LINC00467 stimulates proliferation, migration and invasion capacities of melanoma via targeting miR-485-5p to upregulate PAK1, which provides potential targets for treatment of melanoma.**Keywords:** melanoma, LINC00467, MiR-485-5p, PAK1**Kratak sadržaj****Uvod:** Svrha trenutnog istraživanja je bila da se ispita biološka uloga LINC00467 u izazivanju pogoršanja stanja melanoma.**Metode:** Diferencijalni nivo LINC00467 u tkivima melanoma i njegova prognostička vrednost analizirani su u GEPIA, što je dalje potvrđeno u kliničkim uzorcima koje smo prikupili. Regulatorni efekti LINC00467 na kapacitete proliferacije, migracije i invazije A375 i SKMEL1 ćelijskih linija ispitani su nizom funkcionalnih eksperimenata. Potencijalni nizvodni ciljevi LINC00467 identifikovani su kroz test reportera sa dvostrukom luciferazom, a njihova sinergistička uloga u procesu melanoma je konačno istražena eksperimentima spasavanja.**Rezultati:** LINC00467 je bio pojačano regulisan u uzorcima melanoma, ali nije imao prognostički potencijal kod melanoma. LINC00467 ima kapacitete da stimuliše proliferaciju, migraciju i invaziju A375 i SKMEL1 ćelijskih linija. Identifikovana je povratna sprega LINC00467/miR-485-5p/PAK1, koja je odgovorna za izazivanje pogoršanja melanoma.**Zaključak:** LINC00467 stimuliše kapacitete proliferacije, migracije i invazije melanoma putem ciljanja miR-485-5p da bi se pojačao PAK1, koji pruža potencijalne mete za lečenje melanoma.**Ključne reči:** melanom; LINC00467; MiR-485-5p; PAK1*Address for correspondence:*Yangcheng Li
Address: No.30, Tongyang North Road, Pingchao town,
Nantong, Jiangsu 226000, China
Tel: 86015190821191
e-mail: liyach2010@126.comLixiong Gu
Address: No.20, Xisi Road, Nantong, Jiangsu 226000, China
Tel: 86013962968486
e-mail: ntgulx@163.com

#Zhoujing Ji and Jie Zhang contributed equally to this work

Introduction

Melanoma is a highly aggressive malignancy that develops from uncontrollably proliferated melanocytes. It mainly occurs in young and middle-aged people. Melanoma usually affects skin, but it can also involve the mouth, intestines, or ocular choroid (1, 2). Melanoma is characterized by high degree of malignancy and rapid progression. Current treatment of melanoma is not ideal. Lymphatic or distant metastasis of melanoma develops in the early phase, leading to a poor prognosis with less than 20% of 5-year survival (3, 4). So far, melanoma is still incurable, and its molecular mechanism remains largely unclear.

LncRNAs are over 200 nucleotides long, non-coding RNAs. Although they cannot encode proteins, lncRNAs exert diverse biological functions in the form of RNAs (5, 6). LncRNAs participate in cell functions through transcriptionally, post-transcriptionally or epigenetically regulating gene expressions. They are promising biomarkers for cancer diseases (7–9). Screening melanoma-associated lncRNAs and clarifying their potential functions contribute to develop effective targeted therapy.

We previously analyzed the differential level of LINC00467 in melanoma tissues, which was abnormally up-regulated. In the current study, the potential function of LINC00467 in regulating melanoma cell phenotypes was explored.

Materials and Methods

Melanoma profile

A melanoma profile was analyzed based on the database of GEPIA (<http://gepiacancer-pku.cn/index.html>). Differential level of LINC00467 and its correlation to the survival data in melanoma patients were also analyzed.

Sample collection

Samples of melanoma and paired normal skin tissues were collected from 30 melanoma patients in our hospital from January 2015 to December 2021. Tumor staging was determined based on UICC criteria. None of the enrolled patients received anticancer therapy before surgery that was pathologically proven. Each patient provided informed consent before the study. The present study was approved by the Ethical Committee of Affiliated Hospital of Nantong University.

Cell culture

NHEM and melanoma cell lines (SKMEL1, A375, A2058 and A875) were purchased from ATCC

and were cultivated in Dulbecco's Modified Eagle Medium (DMEM) according to the manufacturer's protocols.

Transfection

Cells grew to 70% density and were then transfected by Lipo 3000. The transfection efficiency of the cells was observed under a microscope 48 hours after transfection. Transfected cells were used for further functional phenotype experiments.

qRT-PCR

RNA from tissue samples is extracted and purified by TRIzol treatment, and the purified RNA is reverse transcribed into cDNAs. Relative expression of PCR products was calculated by $2^{-\Delta\Delta Ct}$ and normalized to that of GAPDH. The primers used were listed as follows: LINC00467: Forward: 5'-ATTGAAGATGCTGCCAAGGG-3'; Reverse: 5'-GCCAGTTTCAGTCCCTCTT-3'; MiR-485-5p: Forward: 5'-CCAAGCTTCACCCATTCCTAACAGGAC-3'; Reverse: 5'-CGGGATCCGTAGGTCAGTTACATGCA-TC-3'; PAK1: Forward: 5'-AAGACATCCAACAGCC-AGAA-3'; Reverse: 5'-TGTAGCCACGTCCCGAGT-3'; GAPDH: Forward: 5'-GGAGCGAGATCCCTCCAAA-AT-3'; Reverse: 5'-GGCTGTTGTCATACTTCTCAT-GG-3'; U6: Forward: 5'-GCTGAGGTGACGGTCTC-AAA-3'; Reverse: 5'-GCCTCCAGTTTCATGGACA-3'.

EdU assay

Cells were treated with 5-ethynyl-2'-deoxyuridine (EdU) for 2 h after cell adhesion, followed by being stained with AdoLo and 4',6-diamidino-2-phenylindole (DAPI). After being washed by PBS for twice, the EdU-labeled cells were observed under a microscope.

Transwell migration assay

A 24-well plate was taken, and 500–600 μ L of culture medium for each group was added to the lower layer, and a Transwell chamber was placed on the top. After digestion, cells were resuspended with serum-free DMEM to determine cell density, and 200 μ L of the desired density of cell suspension was added to each chamber and incubated for 24–72h. Aspirate the culture medium from the upper and lower chambers and add 500 μ L paraformaldehyde (Xavier) to the lower chamber. After 48 h of penetration, cells were fixed, stained, and each sample was captured and counted in five random fields.

Dual-luciferase reporter assay

Binding sequences were predicted using online software for generating luciferase vectors. The Luciferase activity of the seeded cells was determined 48 h after the transfection.

Statistical analysis

SPSS 17.0 (SPSS Inc., Chicago, IL, USA) was adopted for statistical analyses. Survival was evaluated using Kaplan-Meier. $P < 0.05$ indicates statistically significant difference.

Results

LINC00467 was up-regulated in melanoma

Through analyzing melanoma profile downloaded from GEPIA, LINC00467 was up-regulated in melanoma tissues than normal skin tissues (Figure 1A). However, LINC00467 did not have a correlation with the survival data of melanoma (Figure 1B-C). Consistently, in melanoma clinical samples collected in our hospital, LINC00467 was up-regulated (Figure 1D). *In vitro* level of LINC00467 was highly expressed as well (Figure 1E). LINC00467 was of significance in melanoma process.

LINC00467 stimulated proliferative, migratory and invasive functions of melanoma

In A375 and SKMEL1 cells, LINC00467 level was intervened by transfection of oe-LINC00467 or si-LINC00467 (Figure 2A, 2B). EdU assay uncovered that overexpression of LINC00467 enhanced EdU-stained cell ratio in melanoma cells, which was reduced, conversely, by knockdown of LINC00467 (Figure 2C). Both migratory cell rate and invasive cell rate were enhanced in melanoma cells overexpressing LINC00467. Knockdown of LINC00467, on the contrary, reduced metastatic rate (Figure 2D, 2E). Therefore, LINC00467 served as an oncogene in melanoma process.

LINC00467 competitively bound miR-485-5p

The subcellular distribution of LINC00467 was analyzed. The results indicated that LINC00467 was mainly located at the cytoplasm of A375 and SKMEL1 cells, suggesting its post-transcriptional role (Figure 3A). Using Starbase, miR-485-5p was predicted to be a potential target of LINC00467 (Figure 3B). Later, their binding relationship was further verified (Figure 3C). Results showed that miR-485-5p expression was negatively regulated by LINC00467 (Figure 3D). In comparison to normal skin tissues, expression




Figure 1 LINC00467 was up-regulated in melanoma. (A) LINC00467 levels in melanoma profile downloaded from GEPIA; (B) Correlation between LINC00467 and overall survival in melanoma; (C) Correlation between LINC00467 and disease-free survival in melanoma; (D) LINC00467 levels in melanoma tissues and normal ones; (E) LINC00467 levels in melanoma cell lines. * $P < 0.05$; ** $P < 0.01$.




Figure 2 LINC00467 stimulated proliferative, migratory and invasive functions of melanoma. (A) Transfection efficacy of oe-LINC00467 in A375 and SKMEL1 cells; (B) Transfection efficacy of si-LINC00467 in A375 and SKMEL1 cells; (C) EdU-stained A375 and SKMEL1 cells regulated by LINC00467; (D) Migration in A375 and SKMEL1 cells regulated by LINC00467; (E) Invasion in A375 and SKMEL1 cells regulated by LINC00467. *P<0.05; **P<0.01; ***P<0.001.




Figure 3 LINC00467 competitively bound miR-485-5p. (A) Subcellular distribution of LINC00467 in A375 and SKMEL1 cells; (B) Binding sequences in 3'UTR of LINC00467 and miR-485-5p; (C) Direct binding association of LINC00467 with miR-485-5p; (D) MiR-485-5p level in A375 and SKMEL1 cells regulated by LINC00467; (E) MiR-485-5p levels in melanoma tissues and normal ones; (F) A negative association of LINC00467 with miR-485-5p; (G) MiR-485-5p levels in melanoma cell lines. *P<0.05; **P<0.01; ***P<0.001.




Figure 4 PAK1 was the target gene of miR-485-5p. (A) Binding sequences in 3'UTR of miR-485-5p and PAK1; (B) Direct binding relationship between miR-485-5p and PAK1; (C) PAK1 level in A375 and SKMEL1 cells regulated by miR-485-5p; (D) PAK1 levels in melanoma tissues and normal ones; (E) PAK1 levels in melanoma cell lines; (F) A positive correlation between LINC00467 and PAK1; (G) A negative correlation between miR-485-5p and PAK1. * $P<0.05$; ** $P<0.01$.




Figure 5 LINC00467 regulated melanoma cell functions by mediating PAK1. (A) PAK1 level in A375 and SKMEL1 cells co-regulated by PAK1 and LINC00467; (B) EdU-stained A375 and SKMEL1 cells co-regulated by PAK1 and LINC00467; (C) Migration in A375 and SKMEL1 cells co-regulated by PAK1 and LINC00467; (D) Invasion in A375 and SKMEL1 cells co-regulated by PAK1 and LINC00467. ** $P<0.01$ and *** $P<0.001$, vs. NC group; # $P<0.05$ and ## $P<0.01$, vs. oe-PAK1 group.

of miR-485-5p was significantly lower in melanoma tissues (Figure 3E). In addition, the negative association of LINC00467 with miR-485-5p in melanoma tissues was also assessed (Figure 3F). The *in vitro* abundance of miRNA-485-5p was identically down-regulated in melanoma cell lines (Figure 3G).

MiR-485-5p targeted PAK1

Likewise, (P21 (RAC1) Activated Kinase 1) PAK1 was verified as the target gene of miR-485-5p (Figure 4A, 4B). Our findings demonstrated that miR-485-5p negatively regulates PAK1 levels (Figure 4C). QRT-PCR data suggested that level of PAK1 remarkably increased in melanoma cells and tissues (Figure 4D, 4E). It was positively correlated to LINC00467, but negatively correlated to miR-485-5p (Figure 4F, 4G).

LINC00467 regulated melanoma cell functions by mediating PAK1

In A375 and SKMEL1 cells overexpressing PAK1, the up-regulated PAK1 could be inhibited by knockdown of LINC00467 (Figure 5A). Over-expression of PAK1 in melanoma cells remarkably enhanced EdU-stained cell ratio, migratory cell rate and invasive cell rate. However, their enhanced rates were all reversed by knockdown of LINC00467 (Figure 5B–5D). Therefore, PAK1 was responsible for LINC00467-induced melanoma deterioration.

Discussion

Malignant melanoma is derived from melanocytes and it is the most aggressive and deadly skin cancer (10, 11). The global prevalence of melanoma has gradually increased, which seriously endangers human lives and poses a huge economic burden (12). Given above, it's important to explore the molecular mechanism of melanoma.

Previous research have identified the vital function of LINC00467 in several cancers (13–16). Using GEPIA database, LINC00467 was identified to be up-regulated in melanoma tissues, and its up-regulated expression pattern was further confirmed in clinical samples of melanoma and cell lines. Nevertheless, LINC00467 was uncorrelated to the survival data of melanoma. *In vitro* experimentations showed that LINC00467 was capable of stimulating prolifera-

tion, migration and invasion capacities of melanoma cell lines. Since LINC00467 mainly located at the cytoplasm, we believed that it may exert a vital role in post-transcriptional regulation as a ceRNA.

The ceRNA theory proposed that circRNAs, lncRNAs, mRNAs, pseudogenes and other transcripts can competitively bind miRNAs through miRNA response elements (MREs), thus further regulate the downstream targets of miRNAs (17, 18). It is a novel mechanism for explaining RNA interaction. Through bioinformatic prediction and functional analysis, miR-485-5p was discovered to be a possible target of LINC00467.

PAK kinase can be activated by RAS, Rac1 and Cdc42 through GTPase-dependent or non-GTPase-dependent pathways. PAK is widely expressed in tissues, and its activation triggers various life activities. The cancer-associated role of PAK1 has emerged (19–22). Here, PAK1 was found to be responsible for the carcinogenic role of LINC00467 in melanoma. Taken together, a novel feedback loop was identified that LINC00467/miR-485-5p/PAK1 induced melanoma deterioration.

Several limitations should be pointed out. Firstly, only 30 clinical samples of melanoma were collected. More melanoma cases should be recruited in the future to analyze the clinical significance of LINC00467. Secondly, an *in vivo* experiment requires to be designed to validate our findings.

Conclusions

LINC00467 stimulates proliferation, migration and invasion capacities of melanoma through targeting miR-485-5p to upregulate PAK1, which is a possible therapeutic target for melanoma.

Funding Acknowledgements: This study was supported by Nantong Science and Technology Bureau (Peripheral blood dynamic NLR and prognostic nutrition index combined with tumor thickness could guide perioperative treatment and predict prognosis of cutaneous melanoma; Project number: JCZ20211).

Conflict of interest statement

All the authors declare that they have no conflict of interest in this work.

References

1. Brandner JM, Haass NK. Melanoma's connections to the tumour microenvironment. *Pathology* 2013; 45(5): 443–52.
2. Baetz TD, Fletcher GG, Knight G, McWhirter E, Rajagopal S, Song X, et al. Systemic adjuvant therapy for adult patients at high risk for recurrent melanoma: A systematic review. *Cancer Treat Rev* 2020; 87: 102032.
3. Benderli CY. Combination of radiotherapy and vemurafenib in malignant melanoma. *J Buon* 2019; 24(3): 1312–3.
4. Li C, Liu T, Liu B, Hernandez R, Facelli JC, Grossman D. A novel CDKN2A variant (p16(L117P)) in a patient with familial and multiple primary melanomas. *Pigment Cell Melanoma Res* 2019; 32(5): 734–8.
5. Xu S, Gong Y, Yin Y, Xing H, Zhang N. The multiple function of long noncoding RNAs in osteosarcoma progression, drug resistance and prognosis. *Biomed Pharmacother* 2020; 127: 110141.
6. Connerty P, Lock RB, de Bock CE. Long Non-coding RNAs: Major Regulators of Cell Stress in Cancer. *Front Oncol* 2020; 10: 285.
7. Schulte C, Barwari T, Joshi A, Zeller T, Mayr M. Non-coding RNAs versus Protein Biomarkers in Cardiovascular Disease. *Trends Mol Med* 2020; 26(6): 583–96.
8. Huang Z, Zhou JK, Peng Y, He W, Huang C. The role of long noncoding RNAs in hepatocellular carcinoma. *Mol Cancer* 2020; 19(1): 77.
9. Sedano MJ, Harrison AL, Zilaie M, Das C, Choudhari R, Ramos E, et al. Emerging Roles of Estrogen-Regulated Enhancer and Long Non-Coding RNAs. *Int J Mol Sci* 2020; 21(10): 3711.
10. Testori A, Ribero S, Bataille V. Diagnosis and treatment of in-transit melanoma metastases. *Eur J Surg Oncol* 2017; 43(3): 544–60.
11. Hartman ML, Czyn M. MITF in melanoma: mechanisms behind its expression and activity. *Cell Mol Life Sci* 2015; 72(7): 1249–60.
12. Shain AH, Bastian BC. From melanocytes to melanomas. *Nat Rev Cancer* 2016; 16(6): 345–58.
13. Chen Y, Ding Y. LINC00467 enhances head and neck squamous cell carcinoma progression and the epithelial-mesenchymal transition process via miR-299-5p/ubiquitin specific protease-48 axis. *J Gene Med* 2020; 22(7): e3184.
14. Wang H, Guo Q, Nampoukime KB, Yang P, Ma K. Long non-coding RNA LINC00467 drives hepatocellular carcinoma progression via inhibiting NR4A3. *J Cell Mol Med* 2020; 24(7): 3822–36.
15. He X, Li S, Yu B, Kuang G, Wu Y, Zhang M, et al. Up-regulation of LINC00467 promotes the tumorigenesis in colorectal cancer. *J Cancer* 2019; 10(25): 6405–13.
16. Yang J, Liu Y, Mai X, Lu S, Jin L, Tai X. STAT1-induced upregulation of LINC00467 promotes the proliferation migration of lung adenocarcinoma cells by epigenetically silencing DKK1 to activate Wnt/beta-catenin signaling pathway. *Biochem Biophys Res Commun* 2019; 514(1): 118–26.
17. Ding H, Luo Y, Hu K, Liu P, Xiong M. Linc00467 promotes lung adenocarcinoma proliferation via sponging miR-20b-5p to activate CCND1 expression. *Onco Targets Ther* 2019; 12: 6733–43.
18. Li GC, Xin L, Wang YS, Chen Y. Long Intervening Noncoding 00467 RNA Contributes to Tumorigenesis by Acting as a Competing Endogenous RNA against miR-107 in Cervical Cancer Cells. *Am J Pathol* 2019; 189(11): 2293–310.
19. Kim JH, Choi HS, Kim SL, Lee DS. The PAK1-Stat3 Signaling Pathway Activates IL-6 Gene Transcription and Human Breast Cancer Stem Cell Formation. *Cancers (Basel)* 2019; 11(10): 1527.
20. Li X, Zhu J, Liu Y, Duan C, Chang R, Zhang C. MicroRNA-331-3p inhibits epithelial-mesenchymal transition by targeting ErbB2 and VAV2 through the Rac1/PAK1/beta-catenin axis in non-small-cell lung cancer. *Cancer Sci* 2019; 110(6): 1883–96.
21. Zhou Y, Zhang J, Wang J, Cheng MS, Zhao DM, Li F. Targeting PAK1 with the Small Molecule Drug AK963/40708899 Suppresses Gastric Cancer Cell Proliferation and Invasion by Downregulation of PAK1 Activity and PAK1-Related Signaling Pathways. *Anat Rec (Hoboken)* 2019; 302(9): 1571–9.
22. Liang J, Han B, Zhang Y, Yue Q. Numb inhibits cell proliferation, invasion, and epithelial-mesenchymal transition through PAK1/beta-catenin signaling pathway in ovarian cancer. *Onco Targets Ther* 2019; 12: 3223–33.

Received: May 11, 2022

Accepted: September 03, 2022