

# The Effect of Field Dodder (*Cuscuta campestris* Yunck.) on Morphological and Fluorescence Parameters of Giant Ragweed (*Ambrosia trifida* L.)

---

Sava Vrbničanin<sup>1</sup>, Marija Sarić-Krsmanović<sup>2</sup> and Dragana Božić<sup>1</sup>

<sup>1</sup>University of Belgrade, Faculty of Agriculture, Nemanjina 6, 11080 Belgrade, Serbia  
(sava@agrif.bg.ac.rs)

<sup>2</sup>Institute of Pesticides and Environmental Protection, Banatska 31b, 11080 Belgrade, Serbia

Received: February 13, 2013

Accepted: March 14, 2013

## SUMMARY

The effect of the parasitic flowering plant known as field dodder (*Cuscuta campestris* Yunck.) on morphological and fluorescence parameters of infested giant ragweed (*Ambrosia trifida* L.) plants was examined under controlled conditions. The parameters of chlorophyll fluorescence ( $F_o$ ,  $F_v/F_m$ ,  $\Phi_{PSII}$ ,  $F_v$ ,  $F_m$ , ETR and IF) were measured on infested (I) and non-infested (N) *A. trifida* plants over a period of seven days, beginning with the day of infestation. Morphological parameters (plant height, dry and fresh weight) were measured on the last day of fluorescence measurements. *C. campestris* was found to affect the height, fresh and dry weight of the infested *A. trifida* plants, causing significant reduction in plant height and dry weight. Field dodder also affected several parameters of chlorophyll fluorescence ( $F_o$ ,  $F_v/F_m$ ,  $\Phi_{PSII}$  and  $F_v$ ) in infested *A. trifida* plants.

**Keywords:** *Cuscuta campestris*; *Ambrosia trifida*; Morphology; Fluorescence

## INTRODUCTION

*Cuscuta campestris* Yunck. is considered the most widespread plant species of the *Cuscuta* genus. Although North America is assumed to be its place of origin, the species is Cosmopolitan and distributed

throughout South America, Europe, Asia, Africa and Australia (Holm et al., 1997). The genus *Cuscuta* includes predominantly annual and some perennial non-herbaceous plants with threadlike and twining stems, and leaves reduced to inconspicuous scales. As an obligatory stem parasite, *C. campestris* has only a rudimentary root

while its haustoria get attached to the host plant (Swift, 1996); an appressorium is formed at first and then haustoria that penetrate vascular bundles of the host.

Serious damage that field dodder can cause results primarily from the fact that parasitism is the most severe form of negative interaction between vascular plants in which one of the partners moves to heterotrophic nutrition and gets nutrients from the host plant. Dodder-infested plants gradually become weak, lush growth declines and their vegetative and generative yields decrease (Koskela et al., 2001; Fathoulla and Duhoky, 2008). Damage can ultimately lead to total destruction and death of the host. Field dodder causes most damage during massive infestation of recently established leguminous crops (alfalfa or clover) when it challenges the very feasibility of crop production. Problems with field dodder also occur in vegetable nurseries (e.g. tomato, pepper or cabbage), plastic greenhouses, crops such as sugar beet or potato, while hosts of various other species of the *Cuscuta* genus are often weed species, such as *Polygonum aviculare*, *P. persicaria*, *Amaranthus retroflexus*, *Urtica dioica*, *Chenopodium album*, *Cirsium arvense*, *Convolvulus arvensis*, *Ambrosia artemisiifolia*, *Xanthium strumarium*, etc. (Kojić and Vrbničanin, 2000; Rančić and Božić, 2004). The genus *Ambrosia* is a group of invasive weeds and allergenic plants that are widespread across the globe. Species of this genus (e.g. *Ambrosia artemisiifolia*) appeared in this country some 50 years ago and have become a major problem both regarding agricultural purposes and human health as a result of strong allergic reactions that their pollen is able to cause. *Ambrosia trifida* L. is an allochthonous invasive neotophyte, found sporadically in the territory of Serbia (Službeni glasnik R. Srbije, 2010, List IA part II). The latest data have located its foci in central parts of the Bačka region around Kucura, Savino Selo, Ravno Selo and Despotovo, on roadsides in and between villages, as well as around fields and in sunflower, maize, soybean and sugar beet crops (Malidža and Vrbničanin, 2006). During a survey and mapping of allochthonous invasive weeds in Serbia in 2006, the species was not found in the environs of Čoka where Šajinović and Koljadzinski had first detected it in 1982.

As most other *Ambrosia* species, *A. trifida* is also originating in North America (Tamarcaz et al., 2005). It is an annual species exceptionally large in size (exceeding 4 m in height), with branching shoot, large leaf area and weight, allometric sex distribution and high production of large seeds (around 5000 achenes per plant). Considering the damage that *A. trifida* is able to cause, it is necessary to study various measures and develop a strategy

for its control. Herbicides and mechanical control (digging, cutting, ploughing or burning) are the most adequate and most frequently chosen forms of control of this invasive species, as well as some other. However, eradication of invasive weeds by herbicide treatments on non-agricultural areas is too costly, while concern for preserving the surrounding vegetation does not provide much opportunity for herbicide treatments. Besides, chemical control is not acceptable on sites close to water sources, and in gardens and yards around housing facilities. Therefore, a strategy of sustainable control of invasive weeds requires methods for their long-term suppression, including biological control. Biological control has so far included the use of insects, mites, microorganisms, nematodes, herbivorous fish, etc. (Petanović et al., 2000). Using *Cuscuta* species to control invasive plants is a new approach that has been developing in China over the past years (Yu et al., 2009; Shen et al., 2011). It has been confirmed that the species *C. campestris* is fit to be used as an agent of control of the exotic species *Mikania micrantha* (Yu et al., 2009; Shen et al., 2011), which is listed as one of the world's 100 most invasive species (Lowe et al., 2001). Yu et al. (2009) based their conclusions about possibilities for its biological control using *C. campestris* on data for coverage and weight of *M. micrantha* infested by this parasite, and on their analysis of N, P and K contents in infested plants. In a similar study monitoring the same parameters, Shen et al. (2011) also included the effect of *C. campestris* on photosynthesis, stomatal conductance, transpiration, chlorophyll content and content of soluble proteins.

In the present study, we focused on examining the effect of *C. campestris* on the invasive weed species *A. trifida* in order to assess its potentials for biological control of *A. trifida*.

## MATERIAL AND METHODS

The effect of *C. campestris* on morphological and chlorophyll fluorescence parameters was monitored on infested and non-infested *A. trifida* plants. The latter were grown in plastic pots of 10 cm diameter, each containing two *A. trifida* plants. Several parameters of chlorophyll fluorescence were measured over a period of seven days beginning with the first day of infestation (DI) of host plants. Fluorometer PAM-2100 (Heinz Walz, GmbH, Effeltrich, Germany) was used for all measurements, which were taken at 24 h intervals, and the plants were kept for 2 h in the dark before them. The following parameters were measured directly:

minimal fluorescence ( $F_0$  – fluorescence yield in the absence of actinic (photosynthetic) light, i.e. in its basic, non-excited state); maximal fluorescence of darkness-adapted leaf ( $F_m$ ); ratio of variable and maximal fluorescence ( $F_v/F_m$  – maximal fluorescence emitted by photosystem II after absorbing a quantum of light); electron transport rate (ETR); effective fluorescence yield emitted by photosystem II after absorbing a quantum of light ( $\Phi_{PSII}$ ). Two other parameters were computed: intensity of fluorescence (IF) and variable fluorescence ( $F_v$ ). IF was calculated as a ratio of  $F_t/F_0$ , the  $F_t$  being the fluorescence yield under light at stable state and  $F_v$  a calculated difference between the maximal and minimal fluorescence ( $F_v = F_m - F_0$ ). On the last day of fluorescence measurements, the height and fresh weight of plants were also measured. Dry weight was measured after keeping the plants in a drying chamber for 72 h at 60°C. All parameters were measured in four replications and the trial was repeated twice.

Data were processed by the STATISTICA®8.0 software package and average and standard deviation (SD) values were calculated for each parameter. T-test was used to determine significant differences between the infested and non-infested plants for each of the analysed parameters.

## RESULTS AND DISCUSSION

### The effect of *Cuscuta campestris* on morphological parameters of infested *Ambrosia trifida* plants

*C. campestris* has been shown in earlier studies to have a significant effect on weight, flower production and yield of host plants, as well as on their general physiological state (Deng et al., 2003; Zan et al., 2003). Also, some researchers have concluded that parasitic flowering plants of the genus *Cuscuta*, most especially *C. campestris* Yunck., *C. chinensis* Lam. and *C. australis* R. Br., can significantly reduce their host's growth (Liao et al., 2002; Zan et al., 2003; Zhang et al., 2004). Similar data emerged in our own research as *C. campestris* was found to change the height, fresh weight and dry weight of above-ground stems of *A. trifida* (Table 1). All three morphological parameters (plant height, and fresh and dry weight) had higher values in non-infested (14.23±2.80 cm; 2.15±0.51 g; 0.68±0.20 g) than in infested plants (8.85±1.38 cm; 1.80±0.48 g; 0.47±0.11 g). Statistical analysis of these data detected significant differences ( $p < 0.05$ ) in height and dry weight between

non-infested and infested *A. trifida* plants, while the effect of *C. campestris* on fresh weight was not statistically significant ( $p > 0.05$ ) (Table 2). Jeschke et al. (1994, 1997) had found that *Coleus blumei* and *Lupinus albus* parasitized by *Cuscuta reflexa* had much lower weights than non-infested plants. Similar results were later reported by Shen et al. (2005), who measured for 40-50 days the stem/root weight ratio in *Mickania micrantha* plants non-infested and infested by *C. campestris*, and detected significant differences. Other comparable data had been reported by Watling and Press (1997) after investigating the effect of *Striga hermonthica* and *S. asiatica* on *Sorghum bicolor*.

**Table 1.** The effect of *C. campestris* on morphological parameters of *A. trifida* plants

Parameter	Non-infested <i>A. trifida</i> plants	Infested <i>A. trifida</i> plants
Plant height (cm)	14.23±2.80	8.85±1.38
Plant fresh weight (g)	2.15±0.51	1.80±0.48
Plant dry weight (g)	0.68±0.20	0.47±0.11

**Table 2.** Significant differences in morphological parameters between non-infested and infested *A. trifida* plants


Parameter	Non-infested : Infested
Plant height (cm)	0.001756 **
Plant fresh weight (g)	0.250267 <sup>NZ</sup>
Plant dry weight (g)	0.042742 *

<sup>NZ</sup> - not statistically significant ( $p > 0.05$ ); (0.01 <  $p < 0.05$ )\*; ( $p < 0.01$ )\*\*; t-test


### The effect of *Cuscuta campestris* on fluorescence parameters of infested *Ambrosia trifida* plants

Many researchers have used methods based on chlorophyll fluorescence to monitor the effect of various stress factors on plants, such as water deficit (Duraes et al., 2001), extreme temperatures (Francheboud et al., 1999), high salt concentrations (Moradi and Ismail, 2007) or nitrogen deficit (Duraes et al., 2001), to study changes in photosynthetic processes caused by herbicides (Abbaspoor et al., 2006; Pavlović et al., 2007; Bozic et al., 2010), or to examine the resistance of weeds to photosynthesis-inhibiting herbicides (Korres et al., 2003; Pavlovic et al., 2008) or pathogen infection


(Duraes et al., 2001). However, we have discovered no studies that used chlorophyll fluorescence as an indicator of stress in host plants parasitized by *C. campestris*, and our data (Figures 1-4) show that this parasitic flowering plant has a significant ( $p < 0.05$ ) effect on some fluorescence parameters (Table 3). Our analysis of data on the effect of *C. campestris* on parameters of chlorophyll *a* fluorescence in *A. trifida* plants showed that  $F_v$ ,  $F_v/F_m$ ,  $\Phi_{PSII}$  and  $F_v$  were parameters sensitive to parasitic effect. The other fluorescence parameters ( $F_m$ , ETR and IF) were unaffected by *C. campestris*, and were consequently excluded from further study.




**Figure 1.** Parameter  $F_v/F_m$  in non-infested (N) and infested (I) *A. trifida* plants



**Figure 2.** Parameter  $\Phi_{PSII}$  in non-infested (N) and infested (I) *A. trifida* plants



**Figure 3.** Parameter  $F_o$  in non-infested (N) and infested (I) *A. trifida* plants



**Figure 4.** Parameter  $F_v$  in non-infested (N) and infested (I) *A. trifida* plants

**Table 3.** Significant differences for fluorescence parameters in non-infested and infested *A. trifida* plants

Parameter	Non-infested/Infested plants
$F_v/F_m$	0.043902*
$\Phi_{PSII}$	0.003399**
$F_o$	0.144719 <sup>NZ</sup>
$F_v$	0.804431 <sup>NZ</sup>

<sup>NZ</sup>- not statistically significant ( $p > 0.05$ ); ( $0.01 < p < 0.05$ )\*; ( $p < 0.01$ )\*\*; t-test

The stressful influence of *C. campestris* on *A. trifida* led to reductions in the parameters  $F_v$ ,  $F_v/F_m$  and  $\Phi_{PSII}$ , which is consistent with reports from other studies (Abbaspoor et al., 2006; Riethmuller-Haage et al., 2006) that confirmed lower values of these parameters in plants exposed to stress caused by various factors. Contrarily, *C. campestris* was found to increase  $F_o$  values in host plants, compared to non-infested plants, which is consistent with data showing that  $F_o$  values also increased after stressful activity of herbicides in sunflower plants (Božić, 2010). An analysis of all parameters of fluorescence in *A. trifida* plants ( $F_v/F_m$ ,  $\Phi_{PSII}$ ,  $F_o$ ,  $F_v$ ) recorded during the seven-day trial indicates that  $F_v$  is a more sensitive indicator of stress caused by parasitism of *C. campestris* than  $F_v/F_m$ ,  $F_o$  or  $\Phi_{PSII}$ . Decreasing  $F_v$  values in infested *A. trifida* plants were recorded as early as on the first day of trial. These values were lower than control values throughout the seven-day period, reaching their lowest points on the 3<sup>rd</sup> and 6<sup>th</sup> days of infestation (Figure 4). On the other hand, values of the parameter  $F_o$  in infested plants were higher than control values during the seven days of measurement (Figure 3). The parameters  $F_v/F_m$  and  $\Phi_{PSII}$  had nearly identical values in infested and control plants during the initial two days of measurement, but they decreased in infested plants on the 3<sup>rd</sup> day of infestation and reached the lowest point on the fifth day (Figures 1 and 2). A number

of studies (Klem et al., 2002; Abbaspoor and Streibig, 2005; Abbaspoor et al., 2006) have shown that fluorescence parameters reacted to stress at different speed, depending on a number of factors.

## CONCLUSION

*C. campestris* affects the morphological parameters (height, fresh and dry weight) of infested *A. trifida* plants, reducing significantly their height and dry weight. Also, this parasitic plant affects several parameters of chlorophyll fluorescence [minimal fluorescence ( $F_0$ ), variable/maximal fluorescence ratio ( $F_v/F_m$ ), effective fluorescence yield ( $\Phi_{PSII}$ ) and variable fluorescence ( $F_v$ )], showing that these parameters can be used as indicators of *C. campestris* effect on host plants. The results provide a basis for further testing of *C. campestris* as a potential agent of biological control of the invasive weed species *A. trifida*.

## ACKNOWLEDGEMENT

This research was conducted as part of the project III 46008, financed by the Ministry of Education, Science and Technological Development of the Republic of Serbia.

## REFERENCES

- Abbaspoor, M., & Streibig, J.C. (2005). Clodinafop changes the chlorophyll fluorescence induction curve. *Weed Science*, 53(1), 1-9. doi:10.1614/WS-04-131R
- Abbaspoor, M., Teicher, H.B., & Streibig, J.C. (2006). The effect of root-absorbed PSII inhibitors on Kautsky curve parameters in sugar beet. *Weed Research*, 46(3), 226-235. doi:10.1111/j.1365-3180.2006.00498.x
- Božić, D. (2010). *Reakcije korovskih populacija i hibrida sunčokreta prema herbicidima inhibitorima acetolaktat sintetaze*. Doktorska disertacija. Beograd: Poljoprivredni fakultet.
- Božić, D., Vrbničanin, S., Elezović, I., & Sarić, M. (2010). Chlorophyll fluorescence in vivo as the indicator of sunflower susceptibility to ALS-inhibiting herbicides. In *15th EWRS Symposium*, Kaposvar-Hungary. 299.
- Deng, X., Feng, H.L., Ye, W.H., Yang, Q.H., Xu, K.Y., & Cao, H.L. (2003). A study on the control of exotic weed *Mikania micrantha* by using parasitic *Cuscuta campestris*. *Journal of Tropical and Subtropical Botany*, 11(11), 117-122.
- Duraes, F.O.M., Gama, E.E.G., Magalhaes, P.C., Mariel, I.E., Casela, C.R., Oliveira, A.C., Luchiari Junior, A., & Shanahan, J. F. (2002). The usefulness of chlorophyll fluorescence in screening for disease resistance, water stress tolerance, aluminium toxicity tolerance, and N use efficiency in maize. In *Seventh Eastern and Southern Africa Regional Maize Conference*. 356-360.
- Fathoulla, C.N., & Duhoky, M.M.S. (2008). Biological and anatomical study of different *Cuscuta* species. *Kurdistan 1<sup>st</sup> Conference on Biological Sciences*. J Dohuk Univ, 11(1), 22-39.
- Francheboud, Y., Haldimann, P., Leipner, J., & Stamp, P. (1999). Chlorophyll fluorescence as a selection tool for cold tolerance of photosynthesis in maize (*Zea mays* L.). *Journal of Experimental Botany*, 50, 1533-1540.
- Holm, L., Doll, J., Holm, E., Pancho, J., & Herberger, J. (1997). *World Weeds: Natural Histories and Distribution*. (p. 1129). New York: J. Wiley.
- Jeschke, W.D., & Hilpert, A. (1997). Sink-stimulated photosynthesis and sink-dependent increase in nitrate uptake: nitrogen and carbon relations of the parasitic association *Cuscuta reflexa-Ricinus communis*. *Plant, Cell & Environment*, 20, 47-56.
- Jeschke, W.D., Rath, N., Baumel, P., Czygan, F., & Proksch, P. (1994). Modeling flow and partitioning of carbon and nitrogen in the holoparasite *Cuscuta reflexa* Roxb. and its host *Lupinus albus* L. I. Flows between and within the parasitized host. *Journal of Experimental Botany*, 45, 801-812.
- Klem, K., Spundova, M., Hrabalova, H., Naus, J., Vanova, M., Masojidek, J., & Tomek, P. (2002). Comparison of chlorophyll fluorescence and whole-plant bioassays of isotoproturon. *Weed Research*, 42(5), 335-341. doi:10.1046/j.1365-3180.2002.00293.x
- Kojić, M., & Vrbničanin, S. (2000). Parazitski korovi - Osnovne karakteristike, taksonomija, biodiverzitet i rasprostranjenje, I- vilina kosica (*Cuscuta* L.). *Acta herbologica*, 9, 21-28.
- Korres, N.E., Froud-Williams, R.J., & Moss, S.R. (2003). Chlorophyll fluorescence technique as a rapid diagnostic test of the effects of the photosynthetic inhibitor chlorotoluron on two winter wheat cultivars. *Annals of Applied Biology*, 143(1), 53-56. doi:10.1111/j.1744-7348.2003.tb00268.x
- Koskela, T., Salonen, V., & Mutikainen, P. (2001). Interaction of a host plant and its holoparasite: effects of previous selection by the parasite. *Journal of Evolutionary Biology*, 14(6), 910-917. doi:10.1046/j.1420-9101.2001.00352.x
- Liao, W.B., Fan, Q., Wang, B.X., Wang, Y.J., & Zhou, X.Y. (2002). Discovery of three species of *Cuscuta* harming *Mikania micrantha* in South China and their taxonomical identification. *Acta Scientiarum Naturalium Universitatis Sunyatseni*, 41, 54-56.
- Lowe, S., Browne, M., Boudjelas, S., & Poorter, M.D. (2001). *100 of the world's worst invasive alien species*. A selection from the global invasive species database. Auckland, New Zealand: IUCN/SSC Invasive Species Specialist Group (ISSG).

- Malidža, G., & Vrbničanin, S. (2006). Novo nalazište alohtone korovske vrste *Ambrosia trifida* L. na području Vojvodine. In *VIII savetovanje o zaštiti bilja, Zbornik rezimea*, Zlatibor. 44-45.
- Moradi, F., & Ismail, A.M. (2007). Responses of Photosynthesis, Chlorophyll Fluorescence and ROS-Scavenging Systems to Salt Stress During Seedling and Reproductive Stages in Rice. *Annals of Botany*, 99(6), 1161-1173. doi:10.1093/aob/mcm052
- Pavlovic, D., Vrbnicanin, S., Bozic, D., & Fischer, A.J. (2008). Morphophysiological traits and atrazine sensitivity in *Chenopodium album* L. *Pest Manag Sci*, 64(2), 101-7. pmid:18022828. doi:10.1002/ps.1473
- Pavlović, D., Vrbničanin, S., Božić, D., & Simončić, A. (2007). Abutilon theophrasti Medic. Population Responses to Atrazine. *Journal Central European Agriculture*, 8(4), 435-442.
- Petanović, R., Klokočar-Šmit, Z., & Spasić, R. (2000). Biološka borba protiv korova I. *Acta herbologica*, 9, 5-19.
- Rančić, D., & Božić, D. (2004). Uticaj viline kosice (*Cuscuta* sp.) na ptičiji dvornik (*Polygonum aviculare* L.). *Acta herbologica*, 13(1), 167-172.
- Riethmuller-Haage, I., Bastiaans, L., Harbinson, J., Kempenaar, C., & Kropff, M.J. (2006). Influence of the acetolactate synthase inhibitor metsulfuron-methyl on the operation, regulation and organisation of photosynthesis in *Solanum nigrum*. *Photosynthesis Research*, 88(3), 331-341. doi:10.1007/s11120-006-9062-z
- Shen, H., Ye, W.H., Hong, L., Cao, H.L., & Wang, Z.M. (2005). Influence of the obligate parasite *Cuscuta campestris* on growth and biomass allocation of its host *Mikania micrantha*. *Journal of Experimental Botany*, 56, 1277-1284.
- Službeni glasnik Republike Srbije (2010). *Pravilnik o listama štetnih organizama i listama bilja, biljnih proizvoda i propisanih objekata*. broj 7/10 od 19. februara. Retrieved from <http://www.mpt.gov.rs>
- Swift, C. (1996). *Cuscuta* and *Gramica* species dodder a plant parasite.. USA: Colorado State University Cooperative Extension. Tamarcaz, P., Lambelet, C., Clot, B., Keimer, C., & Hauser, C. (2005). Ragweed (*Ambrosia*) progression and its health risks: Will Switzerland resist this invasion. *Swiss Medical Weekly*, 135, 538-548.
- Watling, J.R., & Press, M.C. (1997). Press, M. C. How is the relationship between the C4 cereal *Sorghum bicolor* and the C3 root hemi-parasites *Striga hermonthica* and *Striga asiatica* affected by elevated CO2. *Plant, Cell & Environment*, 20, 1292-1300.
- Zan, Q.J., Wang, B.S., Wang, Y.J., Zhang, J.L., Liao, W.B., & Li, M.G. (2003). The harm caused by *Mikania micrantha* and its control by *Cuscuta campestris*. *Journal of Plant Ecology*, 27, 822-828.
- Zhang, L.Y., Ye, W.H., Cao, H.L., & Feng, H.L. (2004). *Mikania micrantha* H. B. K. in China-an overview. *Weed Research*, 44, 42-49.

## Uticaj viline kosice (*Cuscuta campestris* Yunck.) na morfološke i parametre fluorescencije kod ambrozije trolisne (*Ambrosia trifida* L.)

### REZIME

Ispitivan je uticaj parazitske cvetnice viline kosice (*Cuscuta campestris* Yunck.) na morfološke i parametre fluorescencije kod zaraženih biljaka ambrozije trolisne (*Ambrosia trifida* L.) u kontrolisanim uslovima. Parametri fluorescencije (Fo, Fv/Fm, ΦPSII, Fv, Fm, ETR i IF) kod zaraženih (Z) i nezaraženih (N) biljaka *A. trifida* su mereni sedam dana, počev od prvog dana nakon zaražavanja. Morfološki parametri (visina biljaka, suva i sveža masa biljaka) su mereni poslednjeg dana merenja fluorescencije. *C. campestris* je uticala na visinu, svežu i suhu masu zaraženih biljaka *A. trifida*, pri čemu je dovela do značajne redukcije visine i suve mase biljaka. Takođe, vilina kosica je uticala na veći broj parametara fluorescencije hlorofila (Fo, Fv/Fm, ΦPSII i Fv) kod zaraženih biljaka *A. trifida*.

**Ključne reči:** *Cuscuta campestris*; *Ambrosia trifida*; morfologija; fluorescencija