



Effectiveness of Exercise and Sex Interaction on Adipokines in Obese Rats

Made Mahaguna Putra,^{1, 2} Muchsin Doewes,³ Bambang Purwanto,³ Dono Indarto,³ Hasta Handayani Idrus^{4, 5}

Abstract

Background/Aim: Obesity is a low-grade chronic inflammatory state that promotes a pro-inflammatory immune response. Exercise is considered to be able to prevent inflammation in obesity. The main objective was to determine the effect of the interaction between exercise and sex on inflammation and adipokine in obese rats.

Methods: This research is an experiment using a factorial design with a post-test only. The subjects of this study were 30 Sprague Dawley rats. The rats were 4 months old and weighed 120 g. The rats were given a diet high in carbohydrates and fats. Lee's obesity index was used to measure rats' obesity. The obese rats were then randomly divided into six groups to be given exercise interventions. The intervention provided consisted of aerobic exercise and anaerobic exercise. The primary outcome is adiponectin, leptin, dipeptidyl peptidase-4 (DPP4), tumour necrosis factor alpha (TNF- α) and adipose tissue hypertrophy.

Results: The study found that increased serum adiponectin levels were influenced by exercise ($p < 0.05$) and the interaction between exercise and gender ($p < 0.05$). Decreased serum leptin levels were influenced by gender ($p < 0.05$) and exercise ($p < 0.05$). Decreased serum DPP4 levels were influenced by exercise ($p < 0.05$). Decreased TNF- α expression was influenced by gender ($p < 0.05$) and the interaction between gender and exercise ($p < 0.05$). Decreased adipose cell size was influenced by gender ($p < 0.05$).

Conclusion: This study found that aerobic exercise, anaerobic exercise and the interaction between sex and exercise significantly increased serum adiponectin and decreased TNF- α expression.

Key words: Obesity; Inflammation; Exercise; Adipokines; Sex.

1. Doctoral Student of Medical Sciences, Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Indonesia.
2. Department of Nursing, Sekolah Tinggi Ilmu Kesehatan Rustida, Banyuwangi, East Java, Indonesia.
3. Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Indonesia.
4. Department of Biomedical Sciences, Postgraduate School, Hasanuddin University, Makassar, Indonesia.
5. Centre for Biomedical Research, Research Organisation for Health, National Research and Innovation Agency (BRIN), Jakarta, Indonesia.

Citation: —

Putra MM, Doewes M, Purwanto B, Indarto D, Idrus HH. Effectiveness of exercise and sex interaction on adipokines in obese rats. Scr Med. 2025 Sep-Oct;56(5):935-42.

Corresponding author:

MADE MAHAGUNA PUTRA
E: md.mahagunaputra@gmail.com

Received: 1 February 2025

Revision received: 25 March 2025

Accepted: 25 March 2025

Introduction

Energy imbalance between food intake and energy expenditure will lead to obesity. Low-grade chronic inflammation is associated with obesity, providing signals promoting pro-inflammatory and persistent immunity.^{1, 2} Adipose tissue plays an essential role in regulating obesity.³ In obesity, the size of adipocytes increases (hypertrophy),

consequently increasing macrophage infiltration and ultimately increasing the secretion of various inflammatory cytokines, such as tumour necrosis factor alpha (TNF- α).⁴ TNF- α is a cytokine that regulates glucose and lipid metabolism and increases its concentration in obese patients.⁵

White adipose tissue (WAT) is the primary place for storing excess energy in animals with triacylglycerol stored in adipocytes. WAT targets various hormones and adipokines produced by active endocrine organs, namely polypeptides that act on multiple organs and tissues (muscle, liver and hypothalamus) for general physiological and metabolic regulation. In obesity, several adipokines, such as adiponectin levels, decrease while leptin production increases.⁶ White adipose tissue produces and secretes leptin, whereas peripheral metabolic tissues and the brain secrete leptin receptors. Leptin is essential in appetite suppression and increased energy expenditure, a core component in obesity prevention.⁷ Dipeptidyl peptidase 4 (DPP4) can affect adipogenesis, adipocyte inflammation and insulin sensitivity.⁸ The plasma activity of DPP4 increases with obesity. The study's results explain that adipose tissue releases proinflammatory adipokines, including circulating soluble DPP-4 (sDPP4).⁹

Chronic low-grade inflammation can be reduced by reducing body fat, which increases the production of anti-inflammatory cytokines such as adiponectin. Aerobic exercise is an intervention that can reduce body fat and has anti-inflammatory properties,¹⁰ protects against diseases associated with chronic low-grade inflammation in obesity¹¹ and aerobic exercise can lower adipocyte size and lipid content, resulting in a decrease in adiposity.¹² Anaerobic exercise causes the body to burn more calories and improve insulin sensitivity for hours after rest. It can effectively reduce fat and insulin sensitivity in the body, affect cytokine regulation and maintain a balanced antioxidant status.¹³ Aerobic and anaerobic exercise has extraordinarily beneficial non-pharmacological effects on various functional systems of the human body.¹⁴ Although the beneficial mechanisms of aerobic and anaerobic exercise are unknown, several biological mechanisms have been studied, namely decreased visceral adiposity, triglycerides and low-density lipoproteins,¹⁵ increased high-density lipoproteins,^{16, 17} decreased blood pressure,¹⁸ decreased systemic inflammation¹⁹ and increased insulin sensitivity.²⁰

Based on the background explanation above, researchers are interested in researching the interaction effects of physical exercise and sex on obese rats. This study hypothesises that there is an effect of exercise (aerobic and anaerobic)

interaction with sex on the decrease in TNF- α , increase in adiponectin, decrease in leptin, DPP4 and hypertrophy of adipose tissue in obese model rats. The study aimed to determine what effects aerobic exercise, anaerobic exercise and sex could have on adipokines in obese rats.

Methods

Design

This research was an experiment using a factorial design 2x2 with a post-test only.^{21, 22} This research was conducted at the Integrated Biomedical Laboratory of the Faculty of Medicine, Udayana University. Male and female Sprague-Dawley rats aged 4 months and weighing 120 g were included in this study. Rats were placed in a temperature-controlled room (20-22 °C). The method of calculating the sample size was based on the ANOVA calculation and had an "E" value called the degree of freedom, between 10 and 20. If the value of E was less than 10, then the number of animals was increased and if the value of E was more than 20, then there were enough animals for the experiment. Any sample size with an E value between 10 and 20 was sufficient. The following equation can calculate the value of E: $E = \text{Number of animals} + \text{Number of groups}.$ ²³ Based on these calculations, the animal sample size in this study was 5.

Obese rats that met the obesity Lee index^{24, 25} were randomly divided into three groups: obese rats given aerobic physical exercise, anaerobic physical exercise and obese rats without physical exercise as a baseline. After further intervention, the rats were sacrificed for adipose tissue and blood.

Preparation of rat obesity model

The experimental animals used in this study were 30 Sprague-Dawley rats (*Rattus norvegicus*) obtained from the Faculty of Medicine, Udayana University. Adult male rats were 4-5 months old. Standard rat feed comprises 26 % protein, 63 % carbohydrates and 11 % fat, but in high-fat feed, the composition changes, with 58 % of energy coming from fat.²⁶ Rats were given a diet high in carbohydrates and fats consisting of 22 % carbohydrates, 58 % fat and 10 % protein obtained from Udayana University. The diet was given *ad libitum*, 1x / day, 30 g/rat, for 4

weeks before treatment and continued until treatment was completed. Obesity criteria in rats was obtained by calculating Lee obesity index = $\sqrt[3]{\text{body weight (g)} \times 10 / \text{Nasoanal length (mm)}}$. Rats were declared obese if the Lee obesity index value was > 0.300 .^{24,25}

Exercise

The research equipment used to maintain experimental animals was a plastic tub measuring 45 x 35.5 x 14.5 cm³, a cage lid made of woven wire measuring 36.5 x 28 x 15.5 cm³, water bottles and trash. Rats swimming exercises were performed using a divided box of acrylic measuring 37 x 30 x 50 cm³ with a water depth of 50 cm. Before swimming training began, rats were adapted for 3-5 minutes to adjust to a water temperature of 33 ± 1 °C. Anaerobic exercise were performed on days 1 and 2, 3 and 4 and 5 and 6, rats swam with loads of 10 %, 13 % and 15 % of body mass. The exercise lasted 8 x 10 seconds for 6 days, with an interval of 40 seconds between swimming.²⁷ In aerobic exercise, the rats swam initially for 10 minutes, then increased their swim time by 10 minutes each week to 60 minutes. Next, exercise time was maintained at 60 minutes until day 30/4 of the week.^{28,29}

Adipokines profile

Forty-eight hours after the last training session, 0.5 mL of venous blood was collected via a retro-orbital vein in all the rats. In these samples, adiponectin, leptin and DPP4 were measured by using an ELISA kit. Adiponectin ELISA Kit, Cat. No E0758Ra. Leptin ELISA Kit, Cat. No E0561Ra. Dipeptidyl Peptidase 4 ELISA Kit, Cat. No E0226Ra (*Bioassay Technology Laboratory, Shanghai, China*).

Tissue preparation and histological examination
After the exercise intervention, the rats were sacrificed using a ketamine xylazine dose of 0.1 mg/200 g body weight. White adipose tissue was isolated from subcutaneous fat, fixed in formalin and embedded in paraffin. Five-micrometre slices were cut using a Leica RM550 rotary microtome (*Leica, Vienna, Austria*). The slices were sketched with Mayer Himalayan and eosin. The slides were examined using an Olympus CX41 microscope and an *OptilabPro* camera using *OptilabPro* software.

Immunohistochemistry

Immunohistochemistry for TNF- α was performed on white adipose tissue using a Kit (Dako

EnVision®+ Dual Link System-HRP (DAB+)). Unstained five-micrometre sections were deparaffinised with xylene and rehydrated in a graded ethanol series. Endogenous peroxidase activity was inhibited by 0.3 % H₂O₂. Slides were microwaved using a heat-induced epitope retrieval technique for 3 minutes in tri-sodium citrate buffer. Slides were incubated overnight with anti-TNF- α polyclonal antibody at a dilution of 1:250.

Data analysis

The Kolmogorov-Smirnov test verified the data's normality for all variables. Two-way variance analysis determined the main effects of the type of physical exercise (aerobic and anaerobic exercise), sex and their interaction on serum and adipose tissue variables. A statistically significant difference was considered to be at the level of 0.05. SPSS software, version 16.0, was used for data analysis.

Results

Table 1 shows the results of weight changes and the Lee index. Obese rats that followed aerobic and anaerobic exercise did not change their weight and the Lee index compared to obese rats that did not follow the exercise. Sex affected weight change ($p = 0.000$) and the Lee index ($p = 0.000$) of obese rats who participated in aerobic and anaerobic exercise and did not exercise. Changes in Lee's index were found to be influenced by gender and exercise interaction ($p = 0.038$).

The results of the study (Table 2) found that aerobic exercise and anaerobic exercise were able to increase serum adiponectin levels, serum leptin levels, TNF- α expression (Figure 1) and decrease in adipose cell size (Figure 2) compared to obese rats who did not exercise. The difference in serum adiponectin levels was influenced by exercise ($p = 0.000$) and the interaction between exercise and sex ($p = 0.002$). The difference in serum leptin levels was influenced by sex ($p = 0.032$) and exercise ($p = 0.000$). The difference in serum levels of DPP4 was influenced by exercise ($p = 0.026$). The difference in TNF- α expression was influenced by sex ($p = 0.000$) and the interaction between sex and exercise ($p = 0.012$). The decrease in adipose cell size was affected by sex ($p = 0.000$).

Table 1: Changes in the body weight and Lee index after exercise (n = 30)

Variables	Group (Mean ± SD)			Two-Way ANOVA p-value		
	Aerobic exercise	Anaerobic exercise	Obese	Sex	Exercise	Interaction Sex*Exercise
Initial body weight	222.80 ± 52.57	222.30 ± 60.83	206.80 ± 40.51	-	-	-
Final body weight	221.10 ± 39.82	217.70 ± 53.10	212.90 ± 40.19	0.000	0.695	0.170
Initial Lee index	3.62 ± 0.58	3.60 ± 0.56	3.43 ± 0.39	-	-	-
Final Lee index	3.37 ± 0.40	3.15 ± 0.68	3.49 ± 0.50	0.000	0.423	0.038

Table 2: Mean comparison of adipokines after exercise (n = 30)

Variables	Group (Mean ± SD)			Two-Way ANOVA p-value		
	Aerobic exercise	Anaerobic exercise	Obese	Sex	Exercise	Interaction Sex*Exercise
Adiponectin (mg/L)	3.72 ± 0.29*	4.01 ± 0.47*	2.61 ± 0.97	0.356	0.000	0.002
Leptin (ng/mL)	1.67 ± 0.21*	2.07 ± 0.46*	2.58 ± 0.42	0.032	0.000	0.112
DPP4 (mg/L)	94.39 ± 13.04	91.17 ± 16.26*	111.29 ± 20.22	0.113	0.026	0.626
TNF- α	21.70 ± 10.72	26.20 ± 6.71	26.80 ± 6.95	0.000	0.091	0.012
Adipose tissue hypertrophy (μm)	261.72 ± 102.19	266.81 ± 93.49	315.47 ± 106.49	0.000	0.086	0.680

* Statistically significant at $p < 0.05$, Two-Way ANOVA post hoc test; DPP4: dipeptidyl peptidase 4; TNF- α : tumour necrosis factor alpha;




Figure 1: Immunohistochemistry (IHC) staining of tumour necrosis factor alpha (TNF- α) adipose tissue. IHC staining was performed on adipose tissue from aerobic exercise group, anaerobic exercise group and obesity group n = 10/group. A: IHC staining of TNF- α adipose tissue of aerobic exercise male rats; B: aerobic exercise female rats; C: anaerobic exercise male rats; D: anaerobic exercise female rats; E: obese male rats; F: obese female rats.




Figure 2: Haematoxylin and eosin staining of adipose tissue. Representative images of haematoxylin and eosin staining performed on adipose tissue from aerobic exercise group (AE), anaerobic exercise group (AnE) and obese group (OB) $n = 10/\text{group}$. A: the adipose tissue cell size of aerobic exercise male rats; B: aerobic exercise female rats; C: anaerobic exercise male rats; D: anaerobic exercise female rats; E: obese male rats; F: obese female rats.

Discussion

This study found that sex affects the final body weight and final index of obese rats. The interaction of exercise and gender affects the decrease in the level of obesity in rats. Aerobic exercise and anaerobic exercise increase serum adiponectin and decrease serum leptin. A decrease in serum DPP4 was found in obese rats that had done anaerobic exercise. A decrease in serum leptin, a reduction in TNF- α expression and a decrease in adipose tissue were influenced by sex. The study also found an interaction effect between sex and exercise on decreasing serum adiponectin and decreasing TNF- α expression.

Aerobic exercise can reduce TNF- α in obese male rats, while anaerobic exercise can reduce TNF- α . Research by Jiménez-Maldonado et al found the same thing when running exercise in male mice reduced TNF- α .³⁰ Santiago et al found that

resistance training for 8 weeks can reduce TNF- α in 27 women.³¹ A study conducted by Rejeki et al found that aerobic exercise can reduce leptin levels.³² The study found that exercise in obesity decreases leptin and increases leptin resistance.³³ Bharath et al found decreased body weight, decreased waist circumference and low plasma leptin levels in obese individuals who underwent anaerobic and aerobic exercise.³⁴ However, each gender effect study used the same training session.

This study found an interaction between exercise type and gender on adiponectin levels. Sex is important to an inflammatory response. A comparison of eutrophic groups showed that women had a lower inflammatory status than men. Obese men and women both showed a significantly increased inflammatory response

compared to the eutrophic group, with explicit sex dimorphism in inflammatory profiles in adolescent obesity.³⁵ Aerobic exercise performed by obese male rats can increase adiponectin levels, while obese rats that perform anaerobic exercise increase adiponectin levels. Garekani et al found that aerobic running exercise in obese rats can increase adiponectin levels.³⁶ Rejeki et al also studied the effects of anaerobic exercise in obese women and found that it can increase adiponectin levels.³² Exercise is a safe and inexpensive non-pharmacological intervention to interfere with metabolism.^{37,38} Researchers have stated that increased adiponectin is a beneficial effect of exercise on metabolic disorders.³⁹ The type, volume, intensity of exercise and gender influence adiponectin response.^{40,41}

Significant differences in adipose tissue metabolism between women and men have been reported, with gender as a variable considered in obesity and metabolic disorders.⁴² However, the effect of gender on exercise remains unknown.⁴³ Sex differences in the expression of Aquaporin-7 (AQP7) (associated with genetic predisposition to type 2 diabetes) in women's subcutaneous fat in response to exercise.⁴⁴ There was no difference in the results of methyltetrahydrofolate (MTHFR) epigenetic pattern analysis in sex,⁴⁵ confirming that further studies are needed to investigate the effect of sex. Regulation of adipose inflammatory response by exercise found a decrease in the expression of adiponectin and TNF- α in both sexes. Only leptin decreased in women after anaerobic exercise.⁴⁶ A 2018 meta-analysis review showed aerobic exercise increased adiponectin and decreased leptin levels in diabetic adults, but gender comparisons were absent.⁴⁷ In addition, recent research suggests the impact of exercise on inflammatory adipokines may not differ significantly between men and women.⁴⁸

Conclusion

Decreased adipose tissue hypertrophy, decreased TNF- α expression and decreased DPP4 were found in obese rats who exercised compared to obese rats who did not exercise, but these decreases were not statistically significant. Aerobic and anaerobic exercise increases adiponectin and decreases leptin levels in obesity. Further research needs to be conducted to

optimally affect inflammation and adipokine in obesity by considering longer durations and types of exercise.

Ethics

All animal care and experimental procedures were approved by the Health Research Ethics Commission of Dr Moewardi Regional General Hospital, decision No: 845/VI/HREC/2022, dated 23 June 2022.

Acknowledgement

None.

Conflicts of interest

The authors declare that there is no conflict of interest.

Funding

This article received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Data access

The data that support the findings of this study are available from the corresponding author upon reasonable individual request.

Author ORCID numbers

Made Mahaguna Putra (MMP):
0000-0001-7800-646X
Muchsin Doeves (MD):
0000-0003-4799-2353

Bambang Purwanto (BP):
0009-0007-1881-2719
Dono Indarto (DI):
0000-0002-1057-8725
Hasta Handayani Idrus (HHI):
0000-0002-8563-6395

Author contributions

Conceptualisation: MMP, MD, BP, DI
Methodology: MMP, MD, BP, DI
Validation: MD, BP, DI
Formal analysis: MMP, HHI
Investigation: MMP, MD, BP, DI
Writing - original draft: MMP
Writing - review and editing: MD, BP, DI, HHI.

References

1. Cooke AA, Connaughton RM, Lyons CL, McMorrow AM, Roche HM. Fatty acids and chronic low grade inflammation associated with obesity and the metabolic syndrome. *Eur J Pharmacol.* 2016 Aug 15;785:207-14. doi: 10.1016/j.ejphar.2016.04.021.
2. Bairwa SK, Kumari S, Khandelwal N, Kumar Dhaked G, Dhaked S, Bhatt R. Obesity: an important predictor of metabolic syndrome. *Scr Med* 2023 Mar;54(1):81-5. doi: 10.5937/scriptamed54-40389.
3. Coelho M, Oliveira T, Fernandes R. Biochemistry of adipose tissue: an endocrine organ. *Arch Med Sci.* 2013;9(2):191–200. doi: 10.5114/aoms.2013.33181.
4. Ye J. Mechanisms of insulin resistance in obesity. *Front Med China.* 2013;7(1):14–24. doi: 10.1007/s11684-013-0262-6.
5. Wei Y, Chen K, Whaley-Connell AT, Stump CS, Ibdah JA, Sowers JR. Skeletal muscle insulin resistance: role of inflammatory cytokines and reactive oxygen species. *Am J Physiol Regul Integr Comp Physiol.* 2008 Mar;294(3):R673-80. doi: 10.1152/ajpregu.00561.2007.
6. Leal V de O, Mafra D. Adipokines in obesity. *Clin Chim Acta* 2013;419:87–94. doi: 10.1016/j.cca.2013.02.003.
7. Shin SK, Song SE, Oh JU, Hwang M, Cho HW, Bae JH, et al. Orexin A-induced inhibition of leptin expression and secretion in adipocytes reducing plasma leptin levels and hypothalamic leptin resistance. *Pflugers Arch.* 2019 Dec;471(11–12):1407-18. doi: 10.1007/s00424-019-02318-8.
8. Romacho T, Sell H, Indrakusuma I, Roehrborn D, Castañeda TR, Jelenik T, et al. DPP4 deletion in adipose tissue improves hepatic insulin sensitivity in diet-induced obesity. *Am J Physiol Endocrinol Metab.* 2020 May 1;318(5):E590-E9. doi: 10.1152/ajpendo.00323.2019.
9. Varin EM, Mulvihill EE, Beaudry JL, Pujadas G, Fuchs S, Tanti JF, et al. Circulating levels of soluble dipeptidyl peptidase-4 are dissociated from inflammation and induced by enzymatic DPP4 inhibition. *Cell Metab.* 2019 Feb 5;29(2):320-34.e5. doi: 10.1016/j.cmet.2018.10.001.
10. Gleeson M, Bishop NC, Stensel DJ, Lindley MR, Mastana SS, Nimmo MA. The anti-inflammatory effects of exercise: mechanisms and implications for the prevention and treatment of disease. *Nat Rev Immunol.* 2011 Aug 5;11(9):607-15. doi: 10.1038/nri3041.
11. Ghoshal K, Bhattacharyya M. Adiponectin: Probe of the molecular paradigm associating diabetes and obesity. *World J Diabetes.* 2015;6(1):151. doi: 10.4239/WJD.V6.11.151.
12. Stanford KI, Middelbeek RJW, Goodyear LJ. Exercise effects on white adipose tissue: Beiging and metabolic adaptations. *Diabetes.* 2015;64(7):2361–8. doi: 10.2337/db15-0227.
13. Lee MC, Chung YC, Thenaka PC, Wang YW, Lin YL, Kan NW. Effects of different HIIT protocols on exercise performance, metabolic adaptation, and fat loss in middle-aged and older adults with overweight. *Int J Med Sci.* 2024 Jun 24;21(9):1689-700. doi: 10.7150/ijms.96073.
14. Nieman DC, Wentz LM. The compelling link between physical activity and the body's defense system. *J Sport Heal Sci.* 2018;8(3):201. doi: 10.1016/J.JSHS.2018.09.009.
15. Smith GI, Mittendorfer B, Klein S. Metabolically healthy obesity: facts and fantasies. *J Clin Invest.* 2019;129(10):3978. doi: 10.1172/JCI129186.
16. Warburton DER, Nicol CW, Bredin SSD. Health benefits of physical activity: the evidence. *CMAJ* 2006;174(6):801–9. doi: 10.1503/CMAJ.051351.
17. Mitsui R, Fukushima M, Taniguchi A, Nakai Y, Aoyama S, Takahashi Y, et al. Insulin secretory capacity and insulin sensitivity in impaired fasting glucose in Japanese. *J Diabetes Investig.* 2012 Aug 20;3(4):377-83. doi: 10.1111/j.2040-1124.2012.00201.x.
18. Imazu AA, Goessler KF, Casonatto J, Polito MD. The influence of physical training status on postexercise hypotension in patients with hypertension: a cross-sectional study. *Blood Press Monit.* 2017 Aug;22(4):196-201. doi: 10.1097/MBP.0000000000000255.
19. Batista ML Jr, Rosa JC, Lopes RD, Lira FS, Martins E Jr, Yamashita AS, et al. Exercise training changes IL-10/TNF-alpha ratio in the skeletal muscle of post-MI rats. *Cytokine.* 2010 Jan;49(1):102-8. doi: 10.1016/j.cyto.2009.10.007.
20. Tan X, Chapman CD, Cedernaes J, Benedict C. Association between long sleep duration and increased risk of obesity and type 2 diabetes: A review of possible mechanisms. *Sleep Med Rev.* 2018 Aug;40:127-34. doi: 10.1016/j.smrv.2017.11.001.
21. Collins LM, Dziak JJ, Kugler KC, Trail JB. Factorial experiments: efficient tools for evaluation of intervention components. *Am J Prev Med.* 2014 Oct;47(4):498-504. doi: 10.1016/j.amepre.2014.06.021.
22. Shaw R, Festing MF, Peers I, Furlong L. Use of factorial designs to optimize animal experiments and reduce animal use. *ILAR J.* 2002;43(4):223-32. doi: 10.1093/ilar.43.4.223.
23. Ilyas M, Adzim M, Simbak N, Baig A. sample size calculation for animal studies using degree of freedom (e): an easy and statistically defined approach for metabolomics and genetic research. *Curr Trends Biomed Eng Biosci* 2017;10(2):1-2. doi: 10.19080/ctbeb.2017.10.555785.
24. Damasceno DC, Sinzato YK, Bueno A, Dallaqua B, Lima PH, Calderon IM, et al. Metabolic profile and genotoxicity in obese rats exposed to cigarette smoke. *Obesity (Silver Spring).* 2013 Aug;21(8):1596-601. doi: 10.1002/oby.20152.
25. Campos KE, Volpato GT, Calderon IM, Rudge MV, Damasceno DC. Effect of obesity on rat reproduction and on the development of their adult offspring. *Braz J Med Biol Res.* 2008 Feb;41(2):122-5. doi: 10.1590/s0100-879x2008005000001.
26. King AJF. The use of animal models in diabetes research. *Br J Pharmacol.* 2012;166(3):877-94. doi: 10.1111/j.1476-5381.2012.01911.x.
27. Qiao D, Hou L, Liu X. Influence of intermittent anaerobic exercise on mouse physical endurance and antioxidant components. *Br J Sports Med.* 2006;40(3):214–8. doi: 10.1136/bjsm.2005.020099.
28. Li NC, Wei XX, Hu YL, Hou X, Xu H. Aerobic exercise blocks interleukin-6 levels and germ cell apoptosis in obese rats. *Andrologia.* 2018 Mar;50(2). doi: 10.1111/and.12880.

29. Shaodong C, Haihong Z, Manting L, Guohui L, Zhengxiao Z, Y M Z. Research of influence and mechanism of combining exercise with diet control on a model of lipid metabolism rat induced by high fat diet. *Lipids Health Dis.* 2013 Feb 20;12:21. doi: 10.1186/1476-511X-12-21.

30. Jiménez-Maldonado A, Montero S, Lemus M, Cernea-Cortés J, Rodríguez-Hernández A, Mendoza MA, et al. Moderate and high intensity chronic exercise reduces plasma tumor necrosis factor alpha and increases the Langerhans islet area in healthy rats. *J Musculoskeletal Neuronal Interact.* 2019 Sep 1;19(3):354-61. PMID: 31475943.

31. Macêdo Santiago LÂ, Neto LGL, Borges Pereira G, Leite RD, Mostarda CT, de Oliveira Brito Monzani J, et al. Effects of resistance training on immunoinflammatory response, tnf-alpha gene expression, and body composition in elderly women. *J Aging Res.* 2018 Oct 28;2018:1467025. doi: 10.1155/2018/1467025.

32. Rejeki PS, Pranoto A, Rahmanto I, Izzatunnisa N, Yosika GF, Hernaningsih Y, et al. The positive effect of four-week combined aerobic-resistance training on body composition and adipokine levels in obese females. *Sports (Basel).* 2023 Apr 20;11(4):90. doi: 10.3390/sports11040090.

33. Peng J, Yin L, Wang X. Central and peripheral leptin resistance in obesity and improvements of exercise. *Horm Behav.* 2021;133. doi: 10.1016/j.yhbeh.2021.105006.

34. Bharath LP, Choi WW, Cho JM, Skobodzinski AA, Wong A, Sweeney TE, et al. Combined resistance and aerobic exercise training reduces insulin resistance and central adiposity in adolescent girls who are obese: randomized clinical trial. *Eur J Appl Physiol.* 2018 Aug;118(8):1653-60. doi: 10.1007/s00421-018-3898-8.

35. Simoes E, Correia-Lima J, Sardas L, Storti F, Otani TZDS, Vasques DAC, et al. Sex dimorphism in inflammatory response to obesity in childhood. *Int J Obes (Lond).* 2021 Apr;45(4):879-87. doi: 10.1038/s41366-021-00753-1.

36. Garekani ET, Mohebbi H, Kraemer RR, Fathi R. Exercise training intensity/volume affects plasma and tissue adiponectin concentrations in the male rat. *Peptides.* 2011 May;32(5):1008-12. doi: 10.1016/j.peptides.2011.01.027.

37. Amirazodi M, Mehrabi A, Rajizadeh MA, Bejeshk MA, Esmaeilpour K, Daryanoosh F, et al. The effects of combined resveratrol and high intensity interval training on the hippocampus in aged male rats: An investigation into some signaling pathways related to mitochondria. *Iran J Basic Med Sci.* 2022 Feb;25(2):254-62. doi: 10.22038/IJBM.2022.57780.12853.

38. Rajizadeh MA, Sheibani V, Bejeshk MA, Mohtashami Borzadaran F, Saghari H, Esmaeilpour K. The effects of high intensity exercise on learning and memory impairments followed by combination of sleep deprivation and demyelination induced by etidium bromide. *Int J Neurosci.* 2019 Dec;129(12):1166-78. doi: 10.1080/00207454.2019.1640695.

39. Rajizadeh MA, Esmaeilpour K, Haghparast E, Ebrahimi MN, Sheibani V. Voluntary exercise modulates learning & memory and synaptic plasticity impairments in sleep deprived female rats. *Brain Res.* 2020 Feb 15;1729:146598. doi: 10.1016/j.brainres.2019.146598.

40. Liu K, Zhao W, Li C, Tian Y, Wang L, Zhong J, et al. The effects of high-intensity interval training on cognitive performance: a systematic review and meta-analysis. *Sci Rep.* 2024 Dec 30;14(1):32082. doi: 10.1038/s41598-024-83802-9.

41. Rajizadeh MA, Moslemizadeh A, Hosseini MS, Rafiei F, Soltani Z, Khoramipour K. Adiponectin receptor 1 could explain the sex differences in molecular basis of cognitive improvements induced by exercise training in type 2 diabetic rats. *Sci Rep.* 2023 Sep 27;13(1):16267. doi: 10.1038/s41598-023-43519-7.

42. Chang E, Varghese M, Singer K. Gender and sex differences in adipose tissue. *Curr Diab Rep.* 2018;18(9). doi: 10.1007/S11892-018-1031-3.

43. Landen S, Voisin S, Craig JM, McGee SL, Lamon S, Eynon N. Genetic and epigenetic sex-specific adaptations to endurance exercise. *Epigenetics.* 2019 Jun;14(6):523-35. doi: 10.1080/15592294.2019.1603961.

44. Lebeck J, Østergård T, Rojek A, Füchtbauer EM, Lund S, Nielsen S, et al. Gender-specific effect of physical training on AQP7 protein expression in human adipose tissue. *Acta Diabetol.* 2012 Dec;49 Suppl 1:S215-26. doi: 10.1007/s00592-012-0430-1.

45. Diniz TG, Silva AS, Dos Santos Nunes MK, Ribeiro MD, Filho JM, do Nascimento RAF, et al. Physical activity level influences MTHFR gene methylation profile in diabetic patients. *Front Physiol.* 2021 Jan 12;11:618672. doi: 10.3389/fphys.2020.618672.

46. Da Mota GR, Orsatti FL, Delbin MA, Zanesco A. Resistance exercise improves metabolic parameters and changes adipocyte-derived leptin: a comparison between genders in untrained adults. *Mot Rev Educ Física.* 2016;22(3):217-22. doi: 10.1590/S1980-6574201600030014.

47. Becic T, Studenik C, Hoffmann G. Exercise increases adiponectin and reduces leptin levels in prediabetic and diabetic individuals: systematic review and meta-analysis of randomized controlled trials. *Med Sci.* 2018;6(4). doi: 10.3390/MEDSCI6040097.

48. González-Jurado JA, Suárez-Carmona W, López S, Sánchez-Olivier AJ. Changes in lipoinflammation markers in people with obesity after a concurrent training program: a comparison between men and women. *Int J Environ Res Public Health.* 2020 Aug 25;17(17):6168. doi: 10.3390/ijerph17176168.