



Effect of Deuterated Aspirin Treatment of Obese Iraqi Patients With Blood Hyperviscosity Syndrome

Sura Q Al-Kinany,¹ Fatin Fadhel Mohammed Al-Kazazz,¹ Hussein Inayah,¹ Wasan AM Al-Taie^{2,3,4,5}

Abstract

Background/Aim: Hyperviscosity syndrome (HVS) is a predictor of blood flow resistance and blood viscosity, density or thickness. A number of factors that influence blood viscosity have been identified. This study aimed to compare the effects of the deuterated aspirin (aspirin-D) to standard aspirin (aspirin-H) on the prothrombin time (PT) and the normalised international ratio (INR) in obese HVS patients and healthy individuals of normal weight. Haematocrit (HCT) was the primary determinant of blood viscosity and flow resistance.

Methods: The study included 120 Iraqi individuals, aged between 21 and 55 years, divided into 2 groups, 60 patients in an obese HVS group and 60 in a healthy weight control group. Total complete blood count, haemoglobin (Hb), the effects of aspirin-H and aspirin-D on blood viscosity were evaluated by measuring HCT, red blood cell (RBC), PT and INR.

Results: Patients aged 31–41 years with a body mass index (BMI) of 30–39.9 kg/m² constituted the largest proportion of the HVS group (41.66 %), while the same age group represented the highest percentage among healthy controls (40 %). Patients with HVS showed highly significant increases ($p < 0.001$) in anthropometric and haematological parameters, including BMI, Hb, HCT and RBC, along with a significant reduction in PT and INR compared to controls. Aspirin treatment, particularly deuterated aspirin, significantly increased PT and INR values, indicating improved anticoagulant activity.

Conclusion: Aspirin-D was superior to standard aspirin-H in reducing blood viscosity and delaying clotting time in both HVS patients and healthy subjects.

Key words: Aspirin; Blood viscosity; Deuteration; Hydrogen/Deuterium exchange; Hyperviscosity syndrome.

1. Department of Chemistry, College of Science, Mustansiriyah University, Baghdad, Iraq.
2. Department of Biotechnology, School of Arts and Sciences, American University of Ras Al Khaimah, Ras Al Khaimah, United Arab Emirates.
3. Department of General Education, School of e-Education, Hamdan Bin Mohammed Smart University, Dubai, United Arab Emirates.
4. Emirates Chemical Society, Ras Al Khaimah, United Arab Emirates.
5. Emirates Clinical Chemistry Society, Emirates Medical Association, Dubai, United Arab Emirates.


Citation:

Al-Kinany SQ, Al-Kazazz FFM, Inayah H, Al-Taie WAM. Effect of deuterated aspirin treatment of obese Iraqi patients with blood hyperviscosity syndrome. *Scr Med.* 2026 Mar-Apr;57(2):219-30.

Corresponding author:

SURA Q AL-KINANY
E: suraqaqsim@uomustansiriyah.edu.iq

Received: 29 December 2025
Revision received: 20 March 2026
Accepted: 21 March 2026



Graphical abstract

Introduction

Hyperviscosity syndrome (HVS) is a predictor of blood flow resistance and blood viscosity, density, or thickness.¹ A number of factors that influence blood viscosity have been identified, including the deformability of red blood cell (RBC), packed cell volume (haematocrit - HCT) and plasma viscosity. HCT is the major determinant. When HCT rises, as in erythrocytosis, the viscosity of blood can reach 10-fold higher than that of water.² HVS develops in various haematological diseases, such as leukaemia, sickle cell anaemia, Waldenström macroglobulinaemia (WM) and multiple myeloma (MM).³ When blood viscosity rises above the typical threshold of 4.0 cP, HVS appears.⁴

Patients with HVS experience various symptoms due to poor blood flow, containing headache, confusion, blurred vision, dizziness and thrombotic episodes with and without mucocutaneous haemorrhage.³ High blood viscosity also arises as a result of either a deformation in the shape of RBCs or an abnormal increase in blood proteins, RBC, white blood cells (WBC) and platelets,⁵ or may also result from conditions such as dehydration, smoking, polycythaemia vera and high altitude living. Increased erythropoietin hormone synthesis is the cause of the overproduction of RBC.⁶ The kidneys generate the glycoprotein hormone erythropoietin (EPO), which has 165 amino

acids and stimulates the marrow in the bones to create erythrocytes.⁷ EPO is composed of about 60 % protein and 40 % carbohydrate.⁸

Many antithrombotic drugs are used to avoid complications of HVS, including anticoagulants such as warfarin,⁹ dabigatran,¹⁰ rivaroxaban,¹¹ apixaban,¹² edoxaban,¹³ heparin¹⁴ and antiplatelet (anti-aggregation) agents such as aspirin.¹⁵ Low-dose pharmaceutical blood thinners like warfarin sodium come in clathrate or amorphous forms.⁹ Dabigatran etexilate is licensed in Europe and America for stroke prevention and embolism and for the treatment of deep vein thrombosis.¹

Aspirin (ASA) is a nonsteroidal anti-inflammatory medication (NSAID) recognised for its efficacy in alleviating pain and inflammation. Aspirin, introduced in 1899, became the inaugural effective treatment for rheumatic diseases.¹⁶ It is extensively utilised for the prophylaxis and management of atherosclerotic cardiovascular illnesses¹⁵ and arterial thrombotic disorders owing to its anticoagulant qualities and efficacy in diminishing clot formation resulting from platelet aggregation. Aspirin also helps prevent stroke or other blood vessel diseases.¹⁷ It has been observed that aspirin is effective in preventing or reducing the synthesis of thromboxane

A-2 (TXA2) in platelets. Mechanistically, aspirin blocks the formation of thromboxane through acetylsaline situated in the active site of cyclooxygenase-1 (COX-1). Inhibition of COX-1 results in diminished TXA2 synthesis, an enzyme that enhances platelet formation, facilitates platelet aggregation and serves as a positive feedback mechanism during platelet activation.¹⁸

Hydrogen/deuterium (H/D) isotope exchange is a chemical process wherein hydrogen atoms are substituted with deuterium, or the reverse.¹⁹ The reaction attracted significant interest in the 1960s and 1970s²⁰ and, by the end of the century, there was a great demand for D-labelled compounds, mainly for studies of kinetic isotope effects, chemical reaction mechanisms, the determination of enzyme mechanisms and the improvement of certain drugs.²¹

Isotope-labelled compounds are used in many industries, not least in the pharmaceutical sector. The incorporation of isotopes in an organic chemical reaction can affect the number and profile of products formed. A chemical reaction involving an isotope-containing species is subject to an isotope effect, or an increase or decrease in rate. Sigma bonds with a heavier isotope are

shorter and stronger and have a lower zero-point vibrational energy. These properties are especially significant when isotopes are incorporated into drugs. The replacement of one H atom in a molecule with a larger and heavier D atom can affect the metabolic behaviour of a drug. The carbon-D bond is shorter and stronger than carbon-H, slowing metabolic degradation 6–10-fold. Therefore, H/D exchange can extend the half-life of a drug's efficacy while enhancing the patient's tolerance to it.²² The mechanism of H/D exchange for an aromatic ring is shown in Figure 1.

HVS is an index of blood flow resistance and can also be described as the blood viscosity, mass, or thickness. HCT is a measure of the RBC content of the blood and is the major determinant of blood viscosity. Several antithrombotic agents, including aspirin, have been used in HVS treatment. H/D exchange can confer many metabolic benefits to a drug. According to authors' knowledge, all previous HVS research has concentrated on the impacts of aspirin-H and this is the first to investigate the effects of aspirin-D on blood viscosity. Hypothesis was that deuteration of aspirin, as an anticoagulant medicine, would improve the drug pharmacokinetics and reduce its side effects.




Figure 1: The hydrogen/deuterium (H/D) exchange reaction of an aromatic ring

Figure used with permission of author Dr Hussein Inayah Sharhan²²

Methods

Study population

This research was carried out at Al-Imamian Al-Kadhimiyyin City Hospital and Al-Thuraya laboratory from January 2024 to April 2024. The study comprised 120 Iraqi individuals aged between 21 and 55 years, categorised into two groups: a patient group consisting of 60 individuals diagnosed with HVS and obesity and a healthy control group consisting of 60 individuals with no known health issues. Patients were diagnosed based on clinical biochemical laboratory tests. All patients with chronic diseases (eg angina, stroke, cardiovascular disease) and taking anticoagulant medications and those with cancer, were excluded. The research work was conducted in compliance with the ethical principles for medical research and met the requirements of the Helsinki Declaration.

Specimen collection

Venous blood (5 mL) from each individual was drawn and collected in sodium citrate anticoagulant tubes. After collecting the blood sample and placing it on a roller mixer, it was stored in a refrigerator at 4 °C until use. The blood was then divided into 200 microliter aliquots in three 1.5 mL Eppendorf tubes. Blood samples were used for laboratory tests within 24 h of collection to avoid changes in their composition.²³ The blood was kept at 37 °C during analysis.²⁴

Instruments and materials

The instruments used in this study included a centrifuge (Beckman model TJ-6, Germany), a roller mixer (model VM-370B, Abbott, Taiwan), a STart Max auto coagulation analyser (Stago, France), a Cell-Dyn Ruby automatic haematology analyser (Abbott, USA) and an incubator (Yamato, Japan).

The laboratory chemicals used in this study included haemoglobin (Hb), HCT and RBC kits (Cell-Dyn Ruby, USA). The kit for determination of prothrombin time (PT) and international normalised ratio (INR) was obtained from Stago (France). The PT and INR kit includes lyophilised thromboplastin (R1) and aqueous solvent containing calcium (R2). Aspirin-D (acetylsalicylic acid-D) was synthesised using the H/D exchange method of a previous study.²⁵

Bio-clinical parameters

The sick and control persons were subsequently categorised into two subgroups based on their body mass index (BMI), calculated using the formula: $BMI = \text{weight (kg)} / \text{height (m)}^2$.²⁶ Control healthy persons possess a BMI ranging from 19.9 to 24.9 kg/m², whereas HVS sufferers exhibit a BMI of 30 kg/m² or above. Complete blood counts (CBC), encompassing Hb, HCT and RBC levels, were analysed. The total viscosity was assessed utilising the PT and the international normalised ratio (INR) methodologies in plasma. Anthropometric measures were conducted for both groups.

The lipid accumulation product (LAP) was calculated using the formula: $LAP \text{ (male)} = [\text{waist circumference (WC) (cm)} - 65] \times \text{triglycerides (TG) (mmol/L)}$.²⁷

The visceral adiposity index (VAI) was calculated using the formula: $VAI \text{ (male)} = [\text{WC(cm)} / (39.68 + 1.88 \cdot \text{BMI})] \times (\text{TG}/1.03) \times (1.31 / \text{high-density lipoprotein cholesterol - HDL-C})$, with both TG and HDL levels measured in mmol/L.²⁸

The waist-height ratio (WHtR) was calculated using the formula: $WHtR = \text{waist (cm)} / \text{height (cm)}$ and the waist-hip ratio (WHpR) was calculated using the formula: $WHpR = \text{waist (cm)} / \text{hip (cm)}$. The existence or non-existence of a familial history of HVS was also documented.

Comparative effect of aspirin (H/D) on PT and INR

The optimum conditions for aspirin administration, including concentration (0.03 M), temperature (37 °C) and incubation time (10 min), were determined in a previous study still in the publication process. Three sets of Eppendorf tubes containing 200 µL of whole blood were prepared. In the first set, the values of PT and INR were analysed without adding aspirin. In the second and third sets of blood tubes, the effects of aspirin-H and aspirin-D were assessed under the optimum conditions, whereby aspirin-H (0.03 M) and aspirin-D (0.01 M) were added to the blood and the samples were mixed using a rotary mixer. The blood samples in the second set were incubated with aspirin-H for 10 min at 37 °C in a digital incubator, while no incubation was required in the third set with aspirin-D. Then, the samples in the three sets were centrifuged for 15 min at 2500 rpm. Plasma samples (50 µL) were incubated for 50 s, then the PT and INR reagents (R1 + R2, 100 µL) were added and the values of PT and INR were measured in the three sets.

Calculations

The percentage effects of aspirin-H and aspirin-D on PT and INR were determined using the following equations:

$$\% \text{ effect of aspirin-H} = (PT(A) - PT(H))/PT(A) \times 100$$

$$\% \text{ effect of aspirin-D} = (PT(A) - PT(D))/PT(A) \times 100$$

where PT(A), PT(H) and PT(D) are the sample prothrombin times without aspirin, with aspirin-H and with aspirin-D, respectively. Similar equations were used for the corresponding INR(A), INR(H) and INR(D) values.

Results

The participants' demographic characteristics are displayed in Table 1. A non-significant variation ($p > 0.05$) was seen in the mean age of the HVS patients (36.65 ± 9.56) and that of the healthy control group (35.01 ± 9.13). The mean weight was highly significantly higher ($p < 0.001$) in HVS patients (93.41 ± 15.86 kg) than in the control group (75.46 ± 4.74 kg). The average height exhibited a statistically significant variance ($p < 0.01$) among the HVS patients (1.77 ± 0.07 cm) and the controls (1.81 ± 0.06 cm). The mean BMI was also highly significantly higher in HVS patients ($p < 0.001$) (32.90 ± 2.30 kg/m²) in comparison with the controls (23.13 ± 1.09 kg/m²).

Regarding the WHtR, a highly significant increase ($p < 0.001$) was observed in the HVS patients (0.60 ± 0.06) compared to the control group (0.40 ± 0.02) and the difference in WHpR (0.96 ± 0.13

Statistical analysis

The results were statistically analysed utilising the Statistical Package with the Social Sciences (SPSS) on Windows version 26.0 (IBM Corp, Armonk, New York, USA). Values are shown as mean \pm standard deviation (SD); a p-value beyond 0.05 was deemed non-significant, below 0.05 was regarded as significant and below 0.01 was classified as very significant. The t-test for paired samples was employed to compare mean values.

vs 0.81 ± 0.04 ; $p < 0.001$) was equally significant. The VAI and LAP were also significantly higher in the patient group (4.40 ± 2.71 and 115.70 ± 56.76 , respectively) compared to control group (2.82 ± 1.41 and 22.30 ± 14.69 , respectively). A family history of HVS was recorded for 21.66 % of the HVS patients.

The distribution of BMI in patients with HVS (BMI 30–39.9 kg/m²) and the control group (BMI 19.9–24.9 kg/m²) across various age groups (21–55 years) is illustrated in Figure 2. In the group aged 21–31 years, 11 patients had HVS (18.33 % of the BMI distribution) and 21 represented the control group (35 % of the BMI distribution). The group aged 31–41 years represented 25 HVS patients (41.66 %) and 24 control (40 %) and the group ≥ 41 years represented 24 HVS patients (40 %) and 15 control (25 %).

Table 1: Anthropometric indices in hyperviscosity syndrome (HVS) patients and control group

Parameter	C1 Mean \pm SD	G1 Mean \pm SD	p-value
Age	35.01 \pm 9.13	36.65 \pm 9.56	0.330
Weight (kg)	75.46 \pm 4.74	93.41 \pm 15.86	< 0.001**
Height (m)	1.81 \pm 0.06	1.77 \pm 0.07	0.005**
BMI (kg/m ²)	23.13 \pm 1.09	32.90 \pm 2.30	< 0.001**
WHtR (cm/cm)	0.40 \pm 0.02	0.60 \pm 0.06	< 0.001**
WHpR (cm/cm)	0.81 \pm 0.04	0.96 \pm 0.13	< 0.001**
VAI	2.82 \pm 1.41	4.40 \pm 2.71	< 0.001**
LAP	22.30 \pm 14.69	115.70 \pm 56.76	< 0.001**
Family history (yes, no) %	-	21.66, 78.34 %	-

C1: control group of healthy individuals with normal BMI 19.9–24.9 kg/m²; G1: obese HVS patients with BMI ≥ 30 kg/m²; BMI: body mass index; WHtR: waist-height ratio; WHpR: waist-hip ratio; VAI: visceral adiposity index; LAP: lipid accumulation product; VAI (male) = [waist circumference (WC)(cm) / (39.68 + 1.88·BMI)] \times (triglycerides (TG)/1.03) \times (1.31/ high-density lipoprotein cholesterol - HDL-C), (HDL in mmol/L); LAP (male) = (WC(cm) – 65) \times TG, (TG in mmol/L); (*): significant $p < 0.05$; (**): highly significant $p < 0.01$;




Figure 2: Distribution of body mass index (BMI) in hyperviscosity syndrome (HVS) patients and control group according to age

Complete blood counts for both groups are shown in Table 2. A highly significant increase ($p < 0.001$) was observed in the level of HCT in the HVS patient group ($51.03 \pm 1.79\%$) compared to the control group ($43.82 \pm 1.64\%$). The elevation of Hb (g/dL) and RBC ($10^6/\mu\text{L}$) in the patient group (17.4 ± 0.66 and 5.78 ± 0.36 , respectively) was highly significant ($p < 0.001$) compared to the control group (14.96 ± 0.85 and 5.08 ± 0.26 , respectively).

The values of PT and INR were measured in the absence and presence of aspirin-H and aspirin-D (Table 3). The results indicate a highly significant decrease ($p < 0.001$) in the mean plasma PT in the HVS patient group (12.86 ± 0.59 s) compared to the control group (14.23 ± 0.93 s). In the presence of aspirin-H, the mean plasma PT showed significant elevation ($p < 0.05$) in the HVS patient group

(16.24 ± 1.31 s) compared with the control group (15.78 ± 1.11 s). In the presence of aspirin-D, the mean plasma PT exhibited a highly significant elevation (17.01 ± 1.75 s vs 16.28 ± 1.20 s for patient and control groups, respectively; $p < 0.01$). INR values were significantly decreased ($p < 0.01$) in the HVS patient group (0.95 ± 0.13) compared to the control (1.08 ± 0.26). In the presence of aspirin-H, the INR showed significant elevation ($p < 0.05$) in the HVS patient group (2.57 ± 1.35) compared to the control (2.18 ± 0.84). In the presence of aspirin-D, a similar trend was observed (3.39 ± 1.67 vs 2.73 ± 1.03 ; $p < 0.05$).

The impact of aspirin-D on blood viscosity in HVS patients is detailed in Table 4. Prior to adding aspirin-H or aspirin-D, the PT and INR were 12.86 seconds and 0.95, respectively. Upon adding aspirin-H, an increase in PT and INR was observed,

Table 2: Complete blood counts in the obese hyperviscosity syndrome (HVS) patient and control groups

Blood parameters	C1 Mean \pm SD	G1 Mean \pm SD	p-value
Hb (g/dL)	14.96 ± 0.85	17.4 ± 0.66	$< 0.001^{**}$
HCT (%)	43.82 ± 1.64	51.03 ± 1.79	$< 0.001^{**}$
RBC ($10^6/\text{UL}$)	5.08 ± 0.26	5.78 ± 0.36	$< 0.001^{**}$

C1: control group of healthy individuals with normal body mass index (BMI): $19.9\text{--}24.9$ kg/m²; G1: obese HVS patients with BMI ≥ 30 kg/m²; Hb: haemoglobin; HCT: haematocrit; RBC: red blood cell (erythrocytes); (*): significant $p < 0.05$; (**): highly significant $p < 0.01$;

Table 3: Comparison of PT and INR values between obese hyperviscosity syndrome (HVS) patients and control group

Blood parameters	C1 Mean ± SD	G1 Mean ± SD	p-value
PT without aspirin (s)	14.23 ± 0.93	12.86 ± 0.59	< 0.001**
PT with aspirin-H (s)	15.78 ± 1.11	16.24 ± 1.31	0.040*
PT with aspirin-D (s)	16.28 ± 1.20	17.01 ± 1.75	0.007**
INR without aspirin	1.08 ± 0.26	0.95 ± 0.13	0.001**
INR with aspirin-H	2.18 ± 0.84	2.57 ± 1.35	0.040*
INR with aspirin-D	2.73 ± 1.03	3.39 ± 1.67	0.010**

C1: control group of healthy individuals with normal body mass index (BMI): 19.9–24.9 kg/m²; G1: obese HVS patients with BMI ≥ 30 kg/m²; PT: prothrombin time; INR: international normalised ratio; s: seconds; (*): significant p < 0.05; (**): highly significant p < 0.01;

Table 4: Effect of aspirin-H and aspirin-D on PT and INR in obese hyperviscosity syndrome (HVS) patients

G1 group	(A)	(H)	(D)	% effect aspirin-H	% effect aspirin-D
PT (s)	12.86	15.44	17.01	20.0	32.2
INR	0.95	1.99	3.39	109.4	256.8

Obese HVS patients with body mass index (BMI): ≥ 30 kg/m²; PT(A), PT(H), PT(D): prothrombin time without aspirin, with aspirin-H and with aspirin-D; INR(A), INR(H), INR(D): international normalised ratio without aspirin, with aspirin-H and with aspirin-D;

reaching 15.44 seconds and 1.99, respectively, corresponding to a 20 % increase in PT and a 109.4 % increase in INR. Notably, when aspirin-D was added, an even greater increase in PT and INR was noted, reaching 17.01 seconds (32.2 % increase) and 3.39 (256.8 % increase), respectively.

Discussion

In this study, a group of Iraqis with and without obesity and HVS were examined to determine how aspirin-H and aspirin-D affected blood viscosity. BMI is one of the anthropometric metrics that have been proposed to detect and diagnose obesity.²⁹⁻³¹ Obesity is linked to chronic low-grade inflammation and contributes significantly to the development of chronic illnesses such heart disease and stroke.³² Blood composition can be impacted by elevated inflammatory cytokine levels and inflammation can raise the synthesis of fibrinogen and other clotting factors, which raises blood viscosity.³³ A blood plasma protein called fibrinogen is essential to the clotting process. Obese people often have higher levels of fibrinogen, which can make blood more viscous.³⁴

Obesity often leads to dyslipidaemia (abnormal levels of lipids in the blood), including higher levels of cholesterol and triglycerides. These changes can influence blood viscosity directly and indirectly through interactions with other blood components³⁵ and obesity is frequently accompanied by insulin resistance and elevated insulin levels (hyperinsulinemia). Insulin resistance can lead to endothelial dysfunction, which can promote blood clotting and increase blood viscosity.³⁶

The current study is consistent with that of Brun et al, who found an increase in WHpR in patients with HVS.³⁷ Additionally, in another previous study, subjects with a WHpR greater than 0.90 were shown to have higher blood viscosity than those with a WHpR less than 0.90.³⁸ A further anthropometric measurement proposed to identify and diagnose obesity is the WHtR. WHtR is a reliable indicator of the risk of obesity in both sexes because BMI does not give the distribution of total body fat.³⁹ Measures of central adiposity, including waist circumference (WC) as well as WHpR, have supplanted BMI in a number of clinical diagnostic criteria as more accurate markers of cardiovascular risk linked to obesity.⁴⁰ The distribution of fat in the upper

and lower bodies is taken into consideration by WHpR.⁴¹ Children and adolescents are more likely to be obese and to have related metabolic risks, such as cardiovascular diseases, as a result of increased consumption of calorie-dense foods and sedentary lifestyles.⁴² Recently, the WHtR has been proposed as an effective anthropometric indicator for assessing central adiposity associated with cardiovascular and vascular risk factors.⁴³

The results of this investigation are in line with those of Zhu et al,⁴⁴ who found that patients with HVS had higher levels of LAP and VAI than a control group.⁴⁵ VAI and LAP are measures of the amount of fat accumulated in the organs.⁴⁶ An increase in visceral fat formation and abnormalities in lipid metabolism may be caused by insulin resistance, which in turn could contribute to increased VAI and LAP levels.⁴⁷ Dysregulation of adipocyte activity and lipid metabolism may result from chronic inflammation, a feature shared by HVS. Increased visceral fat deposition and altered lipid profiles may be outcomes of this imbalance.⁴⁸ Hormones including cortisol and adipokines may have an impact on lipid levels in HVS⁴⁹ and the buildup of visceral fat and lipids may be accelerated by hormonal abnormalities.⁵⁰

Non-alcoholic fatty liver disease (NAFLD), which affects liver function, affects lipid metabolism and storage and is linked to elevated VAI and LAP levels.⁵⁰ Some lifestyle and nutritional factors, including obesity, insufficient physical exercise and bad eating habits, may exacerbate visceral fat and lipid buildup in patients with HVS.⁴⁸ Furthermore, there may also be a hereditary component that increases the propensity to exhibit abnormal lipid profiles and thus higher VAI and LAP levels, as well as to build visceral fat.⁵¹

The demographic analysis is illustrated in Table 1. Patients between the ages of 31 and 41 represented the largest percentage of patients with HVS (41.66 %) and a BMI between 30 and 39.9 kg/m², whereas patients between the ages of 21 and 31 had the lowest HVS rate of 18.33 %. On the other hand, the group aged 31–41 years represented the largest percentage (40 %) within the control group (BMI = 19.9–24.9 kg/m²), although this proportion was lower than that of the HVS group. According to Figure 2, the control group's lowest proportion (25 %) was observed among those who were at least 41 years of age.

The findings presented in Table 2 are consistent with those of Visza et al.⁵² Whole blood viscosity

is a complex phenomenon determined by many variables associated with RBCs, with haematocrit considered the most important among them.⁵³ The haematocrit value, representing the percentage of blood volume comprised of RBCs, is the most influential intrinsic factor of blood viscosity and is closely associated with haemoglobin concentration.⁵⁴

Results for HGB and RBC in this current study are consistent with those of Alkhaldy et al.⁵⁵ Viscosity primarily depends on the ability of blood cells to deform under circulatory flow.⁵⁶ RBCs represent the majority of blood cellular components, making the greatest contribution to viscosity.⁵⁷ Hyperviscosity can be caused by elevated haemoglobin levels,⁵⁸ which may result from various disorders. Polycythaemia vera, a bone marrow disorder, causes an excess of RBCs, which raises the total haemoglobin and, in turn, the blood's viscosity.⁵⁹

Table 3 shows how aspirin-H and aspirin-D affect blood viscosity in terms both PT and the INR. Presented results on PT without the addition of aspirin are in line with the research done by Roshal et al.⁶⁰ One of the first laboratory tests for the coagulation system was the PT. It measures how long it takes for a fibrin clot to develop in platelet-poor plasma that has been stimulated with high tissue factor (TF) and anionic phospholipids at the ideal calcium concentration.⁶⁰ The PT test is frequently used to track thrombolytic therapy, preventative therapies and medication therapy. It is an essential laboratory technique for evaluating individuals with coagulation problems.⁶¹

Because aspirin permanently suppresses cyclooxygenase-1 (COX-1), less TA2 is produced, which in turn reduces platelet aggregation. However, as an antiplatelet agent (rather than an anticoagulant like warfarin), aspirin primarily affects platelet aggregation, with a more indirect effect on blood viscosity.⁶²

A decreased INR in the HVS patient group was associated with increased HCT levels (Tables 2 and 3).⁶³ In addition, a difference in blood viscosity was noted between the control group (C1) and the patient group (G1) when taking aspirin-H. The differences in plasma INR levels between the obese HVS patient group and the control group were significant when taking standard aspirin-H and aspirin-D. This could be explained by several factors, including HVS, which alters blood flow and microcirculation, leading to changes in coagulation parameters and INR. The mean values of PT and INR are a clear indication that

aspirin-D reduced the viscosity and delayed the clotting time better than aspirin-H (Table 4).

These findings suggest that the addition of either aspirin-H or aspirin-D exerts an effect on PT and INR in patients with HVS. The highest percentage effect was observed in the presence of aspirin-D. Deuterated aspirin may be more effective than standard aspirin in reducing hyperviscosity and improving prothrombin time due to the effect of deuterium exchange on drug metabolism and activity. When deuterium is added, the metabolism of several medications that are broken down via hydrogen-carbon bond scission pathways is slowed down.⁶⁴ As a result of the kinetic isotope effect, a molecule's physical and chemical properties can be altered by the incorporation of deuterium. The carbon-D bond is more resistant to enzyme cleavage than the carbon-H bond. Therefore, the deuterated molecule may undergo slower metabolism²² and the active drug can stay in the bloodstream for longer, increasing its therapeutic effects.⁶⁵ The enzyme cyclooxygenase (COX) is essential for the production of TA₂, an agent that promotes platelet aggregation and aspirin blocks this enzyme irreversibly.⁶⁶ Deuterated aspirin has a longer half-life than regular aspirin, meaning it is active for a longer period in the body and may have longer-lasting effects on blood viscosity and clotting time, as well as a more sustained suppression of platelet function.⁶⁷

Conclusion

The current study established that the high-risk factor for HVS is obesity with a BMI in excess of 30 kg/m². The findings showed that aspirin-H and aspirin-D were effective in decreasing viscosity and prolonging clotting time and that aspirin-D effected these changes to a greater extent. This is due to the higher carbon-deuterium bond strength, requiring more time for metabolism. The incorporation of deuterium into aspirin led to increased PT and INR. Hence, aspirin-D is a better candidate for controlling platelet aggregation than standard aspirin. Results showed that aspirin-D is superior to standard aspirin-H in reducing viscosity and delaying clotting time. The study findings warrant further clinical investigation and pharmaceutical development of aspirin-D.

Ethics

The research received approval from the Scientific and Ethical Committee at Mustansiriyah University, College of Science, Baghdad, under registration number 51, dated 11 January 2024 and by the Training and Human Development Centre at Baghdad Health Directorate Al-Karkh, Ministry of Health, Baghdad, Iraq, under registration number 17863, dated 7 April 2024.

Acknowledgement

The researchers acknowledge all contributors who gave continuous support and encouragement and thank laboratory staff at Mustansiriyah University, College of Science, Chemistry Department, who facilitated the study in their laboratories.

Conflicts of interest

The authors declare that there is no conflict of interest.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Data access

The data that support the findings of this study are available from the corresponding author upon reasonable individual request.

Author ORCID numbers

Sura Q Al-Kinany (SQAK):
0009-0005-5773-5627

Fatin Fadhel Mohammed Al-Kazazz (FFMAK):
0000-0002-1286-2797
Hussein Inayah (HI):
0000-0002-7045-656X
Wasan AM Al Taie (WAAT):
0009-0000-1145-3329

Author contributions

Conceptualisation: SQAK, WAAT
Methodology: SQAK, FFMAK
Validation: SQAK, HI
Formal analysis: FFMAK, HI
Investigation: SQAK, FFMAK
Resources: SQAK, HI
Writing - original draft: SQAK, FFMAK
Writing - review and editing: SQAK, HI, WAAT
Visualisation: SQAK
Supervision: WAAT
Project administration: SQAK
Funding acquisition: WAAT.

References

- Kucukal E, Man Y, Hill A, Liu S, Bode A, An R, et al. Whole blood viscosity and red blood cell adhesion: potential biomarkers for targeted and curative therapies in sickle cell disease. *Am J Haematol.* 2020;95:1246–56. doi: 10.1002/ajh.25933.
- Kim BJ, Lee YS, Zhbanov A, Yang S. A viscometer for simultaneous measurement of whole blood viscosity and its determinants: haematocrit and red blood cell deformability. *Analyst.* 2019;144:3144–57. doi: 10.1039/c8an02135j.
- Weaver A, Rubinstein S, Cornell RF. Hyperviscosity syndrome in paraprotein secreting conditions including Waldenström macroglobulinemia. *Front Oncol.* 2020;10:815. doi: 10.3389/fonc.2020.00815.
- Al Kuraishy HM, Al Gareeb AI, Al Hamash SM, Cavalu S, El Bouseary MM, Sonbol FI, et al. Changes in the blood viscosity in patients with SARS CoV 2 infection. *Front Med (Lausanne).* 2022;9:876017. doi: 10.3389/fmed.2022.876017.
- Gertz MA. Acute hyperviscosity: syndromes and management. *Blood.* 2018;132:1379–85. doi: 10.1182/blood-2018-06-846816.
- Li K, He C. Gastric mucosal lesions in Tibetans with high altitude polycythemia show increased HIF 1A expression and ROS production. *Biomed Res Int.* 2019;2019:6317015. doi: 10.1155/2019/6317015.
- Sultan AR, Al Kazazz FFM, Mohammed AH. Impact of magnesium oxide nanoparticles on erythropoietin hormone levels in sera of patients with anemia accompanied with diabetic kidney disease. *Nano Biomed Eng.* 2020;12:232–40. doi: 10.5101/nbe.v12i3.p232-240.
- Al Kazazz FFM, Jaber SH, Mohammed AH, Abdullah AH, Kadhim MM, Sultan AR. Preparation of magnesium oxide nanoparticles and study its loaded with recombinant human erythropoietin alfa drug. *Nano Biomed Eng.* 2022;14:186–91. doi: 10.5101/nbe.v14i2.p186-191.
- Dimitrokalli E, Fertaki S, Lykouras M, Kokkinos P, Orkoula M, Kontoyannis C. Warfarin sodium stability in oral formulations. *Molecules.* 2021;26:6631. doi: 10.3390/molecules26216631.
- Abdelghani E, Cua CL, Giver J, Rodriguez V. Thrombosis prevention and anticoagulation management in the pediatric patient with congenital heart disease. *Cardiol Ther.* 2021;10:325–48. doi: 10.1007/s40119-021-00228-4.
- Mladentsev DY, Kuznetsova EN, Skvortsova MN, Dashkin RR. Review on synthetic approaches toward rivaroxaban (Xarelto), an anticoagulant drug. *Org Process Res Dev.* 2022;26:2311–29. doi: 10.1021/acs.oprd.2c00188.
- Dimond M, Looby M, Shah B, Sinha SS, Isseh I, Rollins AT, et al. Design and rationale for the direct oral anticoagulant apixaban in left ventricular assist devices (DOAC LVAD) study. *J Card Fail.* 2024;30:819–28. doi: 10.1016/j.cardfail.2023.10.473.
- Kirchhof P, Toennis T, Goette A, Camm AJ, Diener HC, Becher N, et al. Anticoagulation with edoxaban in patients with atrial high rate episodes. *N Engl J Med.* 2023;389:1167–79. doi: 10.1056/NEJMoa2303062.
- Zang L, Zhu H, Wang K, Liu Y, Yu F, Zhao W. Not just anticoagulation—new and old applications of heparin. *Molecules.* 2022;27:6968. doi: 10.3390/molecules27206968.
- Thobani A, Dhindsa DS, DeMoss BD, Raad M, Sandesara PB, Sperling LS, et al. Usefulness of aspirin for primary prevention of atherosclerotic cardiovascular disease. *Am J Cardiol.* 2019;124:1785–9. doi: 10.1016/j.amjcard.2019.08.040.
- Rauf AMR, Rauf SS, Abed NK. Non steroidal anti-inflammatory drugs (NSAIDs): synthesis of ibuprofen, naproxen and nabumetone. *Al Mustansiriyah J Pharm Sci.* 2019;19:19–27. doi: 10.32947/ajps.v19i3.570.
- Corrigan T, O'Malley L, Bailey D, Moseley H, Okaikoi J, Brown T, et al. Changes in the physical and mechanical properties of human blood with sustained prophylactic use of acetylsalicylic acid (aspirin)—a rheological study. *Open J Fluid Dyn.* 2021;11:167–76. doi: 10.4236/ojfd.2021.114010.
- Badria FA. Drug repurposing: hypothesis, molecular aspects and therapeutic applications. *Norderstedt: BoD—Books on Demand;* 2020. ISBN 978 1 83968 522 4.
- James EI, Murphree TA, Vorauer C, Engen JR, Guttman M. Advances in hydrogen/deuterium exchange mass spectrometry and the pursuit of challenging biological systems. *Chem Rev.* 2021;122:7562–623. doi: 10.1021/acs.chemrev.1c00279.
- Kostyukevich Y, Acter T, Zherbker A, Ahmed A, Kim S, Nikolaev E. Hydrogen/deuterium exchange in mass spectrometry. *Mass Spectrom Rev.* 2018;37:811–53. doi: 10.1002/mas.21565.
- Atzrodt J, Derdau V, Kerr WJ, Reid M. Deuterium and tritium labelled compounds: applications in the life sciences. *Angew Chem Int Ed.* 2018;57:1758–84. doi: 10.1002/anie.201704146.

22. Sharhan HI. H/D exchange using metal colloids in synthesis [thesis]. University of Sussex; 2018. Available from: <http://sro.sussex.ac.uk/>.
23. Trejo Soto C, Hernández Machado A. Normalization of blood viscosity according to the haematocrit and the shear rate. *Micromachines (Basel)*. 2022;13:357. doi: 10.3390/mi13030357.
24. Lee CH, Jung KH, Cho DJ, Jeong SK. Effect of warfarin versus aspirin on blood viscosity in cardioembolic stroke with atrial fibrillation: a prospective clinical trial. *BMC Neurol*. 2019;19:1-7. doi: 10.1186/s12883-019-1315-5.
25. Al Kinany SQ, Inayah H, Al Kazazz FFM. A novel approach to the synthesis of acetylsalicylic acid via H/D substitution. *Pharmakeftiki*. 2025;37(2S):99-103. doi: 10.60988/p.v37i2S.152.
26. Abdulhusein SK, Al Kazazz FFM, Rheima AM. The risk factors in serum of peripheral neuropathy diabetic and type 2 diabetes patients. *Iran J War Public Health*. 2022;14:133-8. doi: 10.29252/ijwph.14.2.133.
27. Ebrahimi M, Seyedi SA, Nabipoorashrafi SA, Rabizadeh S, Sarzaeim M, Yadegar A, et al. Lipid accumulation product (LAP) index for the diagnosis of nonalcoholic fatty liver disease (NAFLD): a systematic review and meta analysis. *Lipids Health Dis*. 2023;22:41. doi: 10.1186/s12944-023-01802-6.
28. Okamura T, Hashimoto Y, Hamaguchi M, Obora A, Kojima T, Fukui M. The visceral adiposity index is a predictor of incident nonalcoholic fatty liver disease: a population based longitudinal study. *Clin Res Hepatol Gastroenterol*. 2020;44:375-83. doi: 10.1016/j.clinre.2019.04.002.
29. Ikeue K, Kusakabe T, Yamakage H, Ishii K, Satoh Asahara N. A body shape index is useful for BMI independently identifying Japanese patients with obesity at high risk of cardiovascular disease. *Nutr Metab Cardiovasc Dis*. 2024;34:387-94. doi: 10.1016/j.numecd.2023.09.008.
30. Lam BCC, Koh GCH, Chen C, Wong MTK, Fallows SJ. Comparison of body mass index, body adiposity index, waist circumference, waist to hip ratio and waist to height ratio as predictors of cardiovascular disease risk factors in an adult population in Singapore. *PLoS One*. 2015;10:e0122985. doi: 10.1371/journal.pone.0122985.
31. Zhu K, Hunter M, James A, Lim EM, Cooke BR, Walsh JP. Relationship between visceral adipose tissue and bone mineral density in Australian baby boomers. *Osteoporos Int*. 2020;31:2439-48. doi: 10.1007/s00198-020-05556-0.
32. Bae YU, Kim Y, Lee H, Kim H, Jeon JS, Noh H, et al. Bariatric surgery alters microRNA content of circulating exosomes in patients with obesity. *Obesity (Silver Spring)*. 2019;27:264-71. doi: 10.1002/oby.22379.
33. Luyendyk JP, Schoenecker JG, Flick MJ. The multifaceted role of fibrinogen in tissue injury and inflammation. *Blood*. 2019;133:511-20. doi: 10.1182/blood-2018-07-818211.
34. Kattula S, Byrnes JR, Wolberg AS. Fibrinogen and fibrin in hemostasis and thrombosis. *Arterioscler Thromb Vasc Biol*. 2017;37:e13-e21. doi: 10.1161/ATVBAHA.117.308564.
35. Zhang T, Chen J, Tang X, Luo Q, Xu D, Yu B. Interaction between adipocytes and high density lipoprotein: new insights into the mechanism of obesity induced dyslipidemia and atherosclerosis. *Lipids Health Dis*. 2019;18:1-11. doi: 10.1186/s12944-019-1170-9.
36. Kaur R, Kaur M, Singh J. Endothelial dysfunction and platelet hyperactivity in type 2 diabetes mellitus: molecular insights and therapeutic strategies. *Cardiovasc Diabetol*. 2018;17:1-17. doi: 10.1186/s12933-018-0763-3.
37. Brun JF, Varlet Marie E, Raynaud de Mauverger E, Mercier J. Both overall adiposity and abdominal adiposity increase blood viscosity by separate mechanisms. *Clin Hemorheol Microcirc*. 2011;48:257-63. doi: 10.3233/CH-2011-1418.
38. Fossum E, Hoieggen A, Moan A, Nordby G, Velund TL, Kjeldsen SE. Whole blood viscosity, blood pressure and cardiovascular risk factors in healthy blood donors. *Blood Press*. 1997;6:161-65. doi: 10.3109/08037059709061932.
39. Elagizi A, Kachur S, Carbone S, Lavie CJ, Blair SN. A review of obesity, physical activity, and cardiovascular disease. *Curr Obes Rep*. 2020 Dec;9(4):571-81. doi: 10.1007/s13679-020-00403-z.
40. Gu Z, Li D, He H, Wang J, Hu X, Zhang P, et al. Body mass index, waist circumference, and waist to height ratio for prediction of multiple metabolic risk factors in Chinese elderly population. *Sci Rep*. 2018;8:1-6. doi: 10.1038/s41598-017-18854-1.
41. Borrueal S, Moltó JF, Alpanes M, Fernandez Duran E, Alvarez Blasco F, Luque Ramirez M, et al. Surrogate markers of visceral adiposity in young adults: waist circumference and body mass index are more accurate than waist hip ratio, model of adipose distribution and visceral adiposity index. *PLoS One*. 2014;9:e114112. doi: 10.1371/journal.pone.0114112.
42. Ejtahed HS, Kelishadi R, Qorbani M, Motlagh ME, Hasani Ranjbar S, Angoorani P, et al. Utility of waist circumference to height ratio as a screening tool for generalized and central obesity among Iranian children and adolescents: the CASPIAN V study. *Pediatr Diabetes*. 2019;20:530-37. doi: 10.1111/pedi.12855.
43. Lo K, Huang YQ, Shen G, Huang JY, Liu L, Yu YL, et al. Effects of waist to height ratio, waist circumference, body mass index on the risk of chronic diseases, all cause, cardiovascular and cancer mortality. *Postgrad Med J*. 2021;97:306-11. doi: 10.1136/postgradmedj-2020-137542.
44. Zhu Y, Huang Y, Sun H, Chen L, Yu H, Shi L, et al. Novel anthropometric indicators of visceral obesity predict the severity of hyperlipidemic acute pancreatitis. *Lipids Health Dis*. 2024;23:120. doi: 10.1186/s12944-024-02112-1.
45. Ding YS, Li Y, Zhang XH, Ma RL, Guo H, Ma L, et al. The improved lipid accumulation product is an accurate index for predicting metabolic syndrome in the Xinjiang population. *Biomed Environ Sci*. 2021;34:503-7. doi: 10.3967/bes2021.070.
46. Demirbas N, Kutlu R. Importance of measured body fat, visceral adiposity index, and lipid accumulation product index in predicting cardiometabolic risk factors. *Metab Syndr Relat Disord*. 2021;19:174-79. doi: 10.1089/met.2020.0098.
47. Başar Gökçen B, Akdevelioğlu Y, Canan S, Bozkurt N. Evaluation of the relationship between serum ferritin and insulin resistance and visceral adiposity index (VAI) in women with polycystic ovary syndrome. *Eat Weight Disord*. 2021;26:1581-93. doi: 10.1007/s40519-020-00980-x.

48. Manna P, Jain SK. Obesity, oxidative stress, adipose tissue dysfunction, and the associated health risks: causes and therapeutic strategies. *Metab Syndr Relat Disord.* 2015;13:423–44. doi: 10.1089/met.2015.0095.
49. Kostapanos MS, Florentin M, Elisaf MS, Mikhailidis DP. Hemostatic factors and the metabolic syndrome. *Curr Vasc Pharmacol.* 2013;11:880–905. doi: 10.2174/15701611113116660171.
50. Tagoe EA, Santa SA. The role of mitochondrial dysfunction in metabolic syndrome. In: Amponsah SK, Ofori EK, Pathak YV, Eds. *Current Trends in the Diagnosis and Management of Metabolic Disorders.* CRC Press; 2023. p.60–85. ISBN 9781003384823.
51. Olubamwo OO. Fatty liver index and risk of incident cardiometabolic disease and mortality [dissertation]. Joensuu (FI): Itä Suomen yliopisto; 2023. ISBN 978 952 61 4798 7.
52. Vizza P, Tradigo G, Parrilla M, Guzzi PH, Gnasso A, Veltri P. On blood viscosity and its correlation with biological parameters. In: ICCS 2018. Cham: Springer; 2018. p.347–53. doi: 10.1007/978-3-319-93701-4_26.
53. Joles JA, Willekes Koolschijn N, Koomans HA. Hypoalbuminemia causes high blood viscosity by increasing red cell lysophosphatidylcholine. *Kidney Int.* 1997;52:761–70. doi: 10.1038/ki.1997.393.
54. Sloop G, Pop GA, St Cyr JA. The haematocrit and blood viscosity are modulated to maintain constant wall shear stress in the carotid sinus. *BIOcomplexity.* 2024;2024. doi: 10.5048/BIO C.2024.2.
55. Alkhaldy HY, Awan ZA, Abouzaid AA, Elbahaey HM, Al Amoudi SM, Shehata SF, et al. Effect of altitude on haemoglobin and red blood cell indices in adults in different regions of Saudi Arabia. *Int J Gen Med.* 2022;3559–65. doi: 10.2147/IJGM.S358139.
56. Méndez Mora L, Cabello Fusarés M, Ferré Torres J, Riera Llobet C, Krishnevskaya E, Trejo Soto C, et al. Blood rheological characterization of β thalassemia trait and iron deficiency anemia using front microrheometry. *Front Physiol.* 2021;12:761411. doi: 10.3389/fphys.2021.761411.
57. Pezzella FR, Antonenko K, Sokolova L. Hyperviscosity syndrome. In: Uluduz D, Arsovska A, Eds. *Rare Causes of Stroke: A Handbook.* 2022. p.162. ISBN 9781108821254.
58. Caimi G, Canino B, Lo Presti R, Urso C, Hopps E. Clinical conditions responsible for hyperviscosity and skin ulcers complications. *Clin Hemorheol Microcirc.* 2017;67:25–34. doi: 10.3233/CH-160218.
59. Vogel J, Kiessling I, Heinicke K, Stallmach T, Ossent P, Vogel O, et al. Transgenic mice overexpressing erythropoietin adapt to excessive erythrocytosis by regulating blood viscosity. *Blood.* 2003;102:2278–84. doi: 10.1182/blood-2003-01-0283.
60. Roshal M, Gil MR. Prothrombin time. In: Shaz BH, Hillyer CD, Gil MR, Eds. *Transfusion Medicine and Hemostasis.* Elsevier; 2019. p.773–77. doi: 10.1016/B978-0-12-813726-0.00128-8.
61. Zaidi SRH, Rout P. Interpretation of blood clotting studies and values (PT, PTT, aPTT, INR, anti factor Xa, D dimer). In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024. PMID: NBK60421.
62. Nam KW, Kwon HM, Jeong HY, Park JH, Kwon H, Jeong SM. High triglyceride/HDL cholesterol ratio is associated with silent brain infarcts in a healthy population. *BMC Neurol.* 2019;19:1–8. doi: 10.1186/s12883-019-1373-8.
63. Almarshad HA, Hassan FM. Alterations in blood coagulation and viscosity among young male cigarette smokers of Al Jouf region in Saudi Arabia. *Clin Appl Thromb Hemost.* 2016;22:386–9. doi: 10.1177/1076029614561319.
64. Belete TM. Recent updates on the development of deuterium containing drugs for the treatment of cancer. *Drug Des Devel Ther.* 2022;3465–72. doi: 10.2147/DDDT.S379496.
65. Qureshi MH. Expansion of the anionic amino Cope cascade and deuteration of monomers for the creation of perdeuterated polymers [dissertation]. Tucson (AZ): University of Arizona; 2022. Available from: <http://hdl.handle.net/10150/667258>.
66. Bruno A, Tacconelli S, Contursi A, Ballerini P, Patrignani P. Cyclooxygenases and platelet functions. In: *Advances in Pharmacology.* Academic Press; 2023. p.133–65. doi: 10.1016/bs.apha.2022.12.001.
67. Jia YM, Gu TT, Ji JZ, Tai T, Zhang MR, Huang BB, et al. Aspirin attenuates the bioactivation of and platelet response to vicagrel in mice. *J Cardiovasc Pharmacol.* 2018;72:252–58. doi: 10.1097/JJC.0000000000000622.