

ОРИГИНАЛНИ НАУЧНИ РАДОВИ  
ОРИГИНАЛЬНЫЕ НАУЧНЫЕ СТАТЬИ  
ORIGINAL SCIENTIFIC PAPERS

## FIXED POINT THEOREM IN A PARTIAL b-METRIC SPACE APPLIED TO QUANTUM OPERATIONS

Rakesh Tiwari<sup>a</sup>, Mohammad Saeed Khan<sup>b</sup>,  
Shoba Rani<sup>c</sup>, Nicola Fabiano<sup>d</sup>

<sup>a</sup> Government V. Y. T. Post-Graduate Autonomous College,  
Department of Mathematics, Durg, Chhattisgarh, Republic of India,  
e-mail: rtiwari@govtsciencecollegedurg.ac.in,  
ORCID iD: <https://orcid.org/0000-0002-6112-0116>

<sup>b</sup> Sefako Makgatho Health Sciences University,  
Department of Mathematics and Applied mathematics,  
Ga-Rankuwa, Republic of South Africa,  
e-mail: drsaeed9@gmail.com,  
ORCID iD: <https://orcid.org/0000-0003-0216-241X>

<sup>c</sup> Government V. Y. T. Post-Graduate Autonomous College,  
Department of Mathematics, Durg, Chhattisgarh, Republic of India,  
e-mail: shobharaniy89@gmail.com,  
ORCID iD: <https://orcid.org/0000-0001-9143-5410>

<sup>d</sup> University of Belgrade, "Vinča" Institute of Nuclear Sciences - National  
Institute of the Republic of Serbia, Belgrade, Republic of Serbia,  
e-mail: nicola.fabiano@gmail.com, **corresponding author**,  
ORCID iD: <https://orcid.org/0000-0003-1645-2071>

DOI: 10.5937/vojtehg70-37109; <https://doi.org/10.5937/vojtehg70-37109>

FIELD: Mathematics

ARTICLE TYPE: Original scientific paper

*Abstract:*

*Introduction/purpose:* A fixed point theorem of an order-preserving mapping on a complete partial b-metric space satisfying a contractive condition is constructed.

*Methods:* Extension of the results of Batsari et al.

*Results:* The fidelity of quantum states is used to construct the existence of a fixed quantum state.

*Conclusions:* The fixed quantum state is associated to an order-preserving quantum operation.



*Key words:* partial b-metric space, order-preserving mapping, quantum operation, fidelity of quantum state, Bloch vector.

## Introduction and preliminaries

A partial metric space is a generalized metric space in which each object does not necessarily have a zero distance from itself (Aamri & El Moutawakil, 2002). Another angle of fixed point research emerged with the approach of the Knaster-Tarski fixed point theorem (Knaster, 1928; Tarski, 1955). The idea was first initiated from Knaster and Tarski in 1927 (Knaster, 1928), and later Tarski found some improvement of the work in 1939, which he discussed in some public lectures between 1939 and 1942 (Tarski, 1955, 1949). Finally, in 1955, Tarski (Tarski, 1955) published the comprehensive results together with some applications. A different property of this theorem is that it involves an order relation defined on the space of consideration. Indeed, the order relation serves as an alternative to the continuity and contraction of the mappings as found in the Brouwer (Brouwer, 1911) and Banach (Banach, 1922) fixed point theorems, respectively, see (Tarski, 1955).

After the approach of the Brouwer (Brouwer, 1911), Banach (Banach, 1922) and Knaster-Tarski (Tarski, 1955) fixed point theorems, many researchers become involved in extension (Browder, 1959; Leray & Schauder, 1934; Schauder, 1930), generalization (Batsari et al, 2018; Browder, 1959; Du et al, 2018) and improvements (Batsari et al, 2018; Batsari & Kumam, 2018; Kannan, 1972; Khan et al, 1984) of the theorems using different spaces and functions. In the way of generalizing spaces was Bourbaki-Bakhtin-Cezerwik's b-metric space (Bakhtin, 1989; Bourbaki, 1974; Czerwinski, 1993), Matthews's partial metric space (Matthews, 1994) and Shukla's Partial b-metric space (Shukla, 2014).

In the area of the quantum information theory, a qubit is seen as a quantum system, whereas a quantum operation can be inspected as the measurement of a quantum system; it describes the development of the system through the quantum states. Measurements have some errors which can be corrected through quantum error correction codes. The quantum error correction codes are easily developed through the information-preserving structures with the help of the fixed points set

of the associated quantum operation. Therefore, the study of quantum operations is necessary in the field of the quantum information theory, at least in developing the error correction codes, knowing the state of the system (qubit) and the description of energy dissipation effects due to loss of energy from a quantum system ([Nielsen & Chuang, 2000](#)).

In 1951, Luders ([Lüders, 1950](#)) discussed the compatibility of quantum states in measurements (quantum operations). He also proved that the compatibility of quantum states in measurements is equivalent to the commutativity of the states with each quantum effects in the measurement.

In 1998, Busch et al. ([Busch & Singh, 1998](#)) generalized the Luders theorem. He also showed that a state is unchanged under a quantum operation if the state commutes with every quantum effect that relates the quantum operation. In 2002, Arias et al. ([Arias et al, 2002](#)) studied the fixed point sets of a quantum operation and gave some conditions for which the set is equal to a commute set of the quantum effects that described the quantum operation. In 2011, Long and Zhang ([Zhang & Ji, 2012](#)) deliberated the fixed point set for quantum operations, they presented some necessary and sufficient conditions for the existence of a non-trivial fixed point set. Similarly, in 2012, Zhang and Ji ([Long & Zhang, 2011](#)) deliberated the existence of a non-trivial fixed point set of a generalized quantum operation. In 2016, Zhang and Si ([Zhang & Si, 2016](#)) explored the conditions for which the fixed point set of a quantum operation ( $\phi_{\mathcal{A}}$ ) with respect to a row contraction  $\mathcal{A}$  equals to the fixed point set of the power of the quantum operation ( $\phi_{\mathcal{A}}^j$ ) for some  $1 \leq j < +\infty$ . Other useful references are ([Agarwal et al, 2015](#); [Debnath et al, 2021](#); [Kirk & Shahzad, 2014](#)).

**DEFINITION 1.** ([Shukla, 2014](#)) A **partial b-metric** on the set  $X$  is a function  $p_s : X \times X \rightarrow \mathbb{R}_+$  such that,

- (1) For all  $x, y \in X$ ,  $x = y$  iff  $p_s(x, x) = p_s(x, y) = p_s(y, y)$
  - (2) For all  $x, y \in X$ ,  $p_s(x, x) \leq p_s(x, y)$
  - (3) For all  $x, y \in X$ ,  $p_s(x, y) = p_s(y, x)$
  - (4) There exists a real number  $s \geq 1$  such that, for all  $x, y, z \in X$ ,  $p_s(x, z) \leq s[p_s(x, y) + p_s(y, z)] - p_s(y, y)$ .
- $(X, p_s)$  denotes the partial b-metric space. Note that every partial metric is



a partial b-metric with  $s = 1$ . Also, every b-metric is a partial b-metric with  $p_s(x, x) = 0$ , for all  $x, y \in X$ .

A sequence  $\{x_n\}$  in the space  $(X, p_s)$  converges with respect to the topology  $\tau_b$  to a point  $x \in X$ , if and only if

$$\lim_{n \rightarrow +\infty} p_s(x_n, x) = p_s(x, x). \quad (1)$$

The sequence  $\{x_n\}$  is Cauchy in  $(X, p_s)$  if the below limit exists and is finite

$$\lim_{n,m \rightarrow +\infty} p_s(x_n, x_m) < +\infty. \quad (2)$$

A partial b-metric space  $(X, p_s)$  is complete, if every Cauchy sequence  $\{x_n\}$  in  $(X, p_s)$  converges to a point  $x \in X$  such that,

$$\lim_{n,m \rightarrow +\infty} p_s(x_n, x_m) = p_s(x, x). \quad (3)$$

**DEFINITION 2.** A mapping  $T$  is said to be order-preserving on  $X$ , whenever  $x \preceq y$  implies  $T(x) \preceq T(y)$  for all  $x, y \in X$ .

## Main result

The objective of this work is to establish a fixed point theorem in a complete partial b-metric space.

**THEOREM 1.** Let  $(X, p_s)$  be a complete partial b-metric space with  $s \geq 1$  and associated with a partial order  $\preceq$ . Suppose an order preserving mapping  $T : X \rightarrow X$  satisfies

$$\begin{aligned} p_s(T(x), T(y)) &\leq \alpha \max\{p_s(x, y), p_s(x, T(y)), p_s(y, T(x))\} \\ &+ \frac{\beta}{2} \min\{p_s(x, T(y)) + p_s(y, T(x)), p_s(x, T(x)) + p_s(y, T(y))\} \end{aligned} \quad (4)$$

for all comparable  $x, y \in X$ , where  $\alpha, \beta \in [0, \theta]$  and  $\theta = \min\{\frac{1}{s^3}, \frac{2}{s+1}\}$ . If there exists  $x_0 \in X$  such that  $x_0 \preceq T(x_0)$ , then  $T$  has a unique fixed point  $\hat{x} \in X$  such that  $p_s(\hat{x}, \hat{x}) = 0$ .

*Proof.* Suppose  $x_0 \neq T(x_0)$ , define a sequence  $\{x_n\} \subseteq X$  by  $x_n = T^n(x_0)$  and let  $q_n = p_s(x_n, x_{n+1})$ . It is clear that if  $x_n = x_{n+1}$  for some natural

number  $n$ , then  $x_n$  is a fixed point of  $T$ , i.e.,  $x_{n+1} = T(x_n) = x_n$ . Let  $x_{n+1} \neq x_n$  for all  $n \in N$ . Then, we proceed as follows:

$$\begin{aligned}
q_n &= p_s(x_n, x_{n+1}) = p_s(T(x_{n-1}), T(x_n)) \\
&\leq \alpha \max\{p_s(x_{n-1}, x_n), p_s(x_{n-1}, T(x_n)), p_s(x_n, T(x_{n-1}))\} \\
&\quad + \frac{\beta}{2} \min\{p_s(x_{n-1}, T(x_n)) + \\
&\quad p_s(x_n, T(x_{n-1})), p_s(x_{n-1}, T(x_{n-1})) + p_s(x_n, T(x_n))\} \\
&= \alpha \max\{p_s(x_{n-1}, x_n), p_s(x_{n-1}, x_{n+1}), p_s(x_n, x_n)\} \\
&\quad + \frac{\beta}{2} \min\{p_s(x_{n-1}, x_{n+1}) + p_s(x_n, x_n), p_s(x_{n-1}, x_n) + p_s(x_n, x_{n+1})\} \\
&= \alpha \max\{p_s(x_{n-1}, x_n), s[p_s(x_{n-1}, x_n) + p_s(x_n, x_{n+1})]\} \\
&\quad + \frac{\beta}{2} \left[ \frac{s[p_s(x_{n-1}, x_n) + p_s(x_n, x_{n+1})] + p_s(x_{n-1}, x_n) + p_s(x_n, x_{n+1})}{2} \right] \\
&= \alpha(s[p_s(x_{n-1}, x_n) + p_s(x_n, x_{n+1})]) + \\
&\quad \frac{\beta}{2}(s+1) \left[ \frac{p_s(x_{n-1}, x_n) + p_s(x_n, x_{n+1})}{2} \right] \\
&= \alpha s + \frac{\beta(s+1)}{4} (p_s(x_{n-1}, x_n) + p_s(x_n, x_{n+1})) \\
&= \frac{4\alpha s + \beta s + \beta}{4} (p_s(x_{n-1}, x_n) + p_s(x_n, x_{n+1})) \\
&= \frac{4\alpha s + \beta s + \beta}{4} (q_{n-1} + q_n).
\end{aligned}$$

Thus, we have

$$q_n \leq \frac{4\alpha s + \beta s + \beta}{4} (q_{n-1} + q_n)$$

which implies

$$\left( \frac{4 - 4\alpha s - \beta s - \beta}{4} \right) q_n \leq \left( \frac{4\alpha s + \beta s + \beta}{4} \right) q_{n-1} \quad (5)$$

By simplifying (5), we have

$$q_n \leq \left( \frac{4\alpha s + \beta s + \beta}{4 - 4\alpha s - \beta s - \beta} \right) q_{n-1} \quad (6)$$

For  $\theta \in \min\{\frac{1}{s^3}, \frac{2}{s+1}\}$ , we deduce that

$$0 \leq \frac{4\alpha s + \beta s + \beta}{4 - 4\alpha s - \beta s - \beta} \leq 1.$$



Therefore, from(6), we conclude that  $p_s(x_n, x_{n+1}) = q_n \leq q_{n-1} = p_s(x_{n-1}, x_n)$ . Thus,  $\{q_n\}_{n=1}^{+\infty}$  is a monotone non-increasing sequence of real numbers and bounded below by 0. Therefore,  $\lim_{n \rightarrow +\infty} q_n = 0$ , see Chidume et al.(Chidume & Chidume, 2014).

Next, we show  $\{x_n\}_{n=1}^{+\infty}$  is Cauchy. Let  $x_n, x_m \in X$ , for all  $n, m \in \mathbb{N}$ . Then,

$$\begin{aligned}
 p_s(x_n, x_m) &= p_s(T^n x_0, T^m x_0) \\
 &= p_s(T(x_{n-1}), T(x_{m-1})) \\
 &= \alpha \max\{p_s(x_{n-1}, x_{m-1}), p_s(x_{n-1}, x_m), p_s(x_n, x_{m-1})\} \\
 &\quad + \frac{\beta}{2} \min\{p_s(x_{n-1}, x_m) + p_s(x_{m-1}, x_n), p_s(x_{n-1}, x_n) + p_s(x_{m-1}, x_m)\} \\
 &= \alpha \max\{s(p_s(x_{n-1}, x_n) + p_s(x_n, x_{m-1})), p_s(x_{n-1}, x_m), p_s(x_n, x_{m-1})\} \\
 &\quad + \frac{\beta}{2} \left\{ \frac{p_s(x_{n-1}, x_m) + p_s(x_{m-1}, x_n) + p_s(x_{n-1}, x_n) + p_s(x_{m-1}, x_m)}{2} \right\} \\
 &= \alpha \max\{s(p_s(x_{n-1}, x_n) + s(p_s(x_n, x_m) + p_s(x_m, x_{m-1}))), p_s(x_{n-1}, x_n) \\
 &\quad + p_s(x_{m-1}, x_m)\} + \frac{\beta}{4} (s(p_s(x_{n-1}, x_n) + p_s(x_n, x_m)) + s(p_s(x_{m-1}, x_m) \\
 &\quad + p_s(x_m, x_n)) + p_s(x_{n-1}, x_n) + p_s(x_{m-1}, x_m)) \\
 &= \alpha s(p_s(x_{n-1}, x_n) + s(p_s(x_n, x_m) + p_s(x_m, x_{m-1}))) \\
 &\quad + \frac{\beta}{4} (sp_s(x_{n-1}, x_n) + 2sp_s(x_n, x_m) + sp_s(x_{m-1}, x_m) \\
 &\quad + p_s(x_{n-1}, x_n) + p_s(x_{m-1}, x_m)) \\
 &\leq (\alpha s + \frac{s\beta}{4} + \frac{\beta}{4}) p_s(x_{n-1}, x_n) + (\alpha s^2 + \frac{s\beta}{2}) p_s(x_n, x_m) \\
 &\quad + (\alpha s^2 + \frac{s\beta}{4} + \frac{\beta}{4}) p_s(x_{m-1}, x_m), \tag{7}
 \end{aligned}$$

implies that

$$\begin{aligned}
 (1 - (\alpha s^2 + \frac{s\beta}{2})) p_s(x_n, x_m) &\leq (\alpha s + \frac{s\beta}{4} + \frac{\beta}{4}) p_s(x_{n-1}, x_n) + \\
 &\quad (\alpha s^2 + \frac{s\beta}{4} + \frac{\beta}{4}) p_s(x_{m-1}, x_m) \\
 &\leq \frac{2}{2 - 2\alpha s^2 - s\beta} ((\alpha s + \frac{s\beta}{4} + \frac{\beta}{4}) p_s(x_{n-1}, x_n) \\
 &\quad + (\alpha s^2 + \frac{s\beta}{4} + \frac{\beta}{4}) p_s(x_{m-1}, x_m)). \tag{8}
 \end{aligned}$$

Now, taking the limit as  $n, m \rightarrow +\infty$  in (7), we have

$$\lim_{n,m \rightarrow +\infty} p_s(x_n, x_m) = 0.$$

Therefore,  $\{x_n\}$  is a Cauchy sequence in  $X$ . For  $X$  being complete, there exists  $\hat{x} \in X$  such that

$$\lim_{n \rightarrow +\infty} p_s(x_n, \hat{x}) = \lim_{n,m \rightarrow +\infty} p_s(x_n, x_m) = p_s(\hat{x}, \hat{x}) = 0.$$

Now, we proceed to prove the existence of the fixed point of  $T$  satisfying (1). Let  $x_0 \in X$  be such that  $x_0 \preceq T(x_0)$ . If  $T(x_0) = x_0$  then,  $x_0$  is a fixed point of  $T$ . Recall that,  $T$  is order-preserving and  $x_0 \preceq T(x_0)$  then, we have  $x_0 \preceq T(x_0) = x_1, x_1 \preceq T(x_1) = x_2, x_2 \preceq T(x_2) = x_3, \dots, x_n \preceq T(x_n) = x_{n+1}$ . By transitivity of  $\preceq$ , we have  $x_0 \preceq x_1 \preceq x_2 \preceq x_3 \preceq \dots \preceq x_n \preceq x_{n+1} \preceq \dots$ .

For showing  $\hat{x} \in X$  is a fixed point of  $T$ , we proceed as follows:

$$\begin{aligned} p_s(\hat{x}, T(\hat{x})) &\leq s[p_s(\hat{x}, x_{n+1}) + p_s(x_{n+1}, T(\hat{x})) - p_s(x_{n+1}, x_{n+1})] \\ &\leq s[p_s(\hat{x}, x_{n+1}) + p_s(T(x_n), T(\hat{x}))] \\ &\leq s[p_s(\hat{x}, x_{n+1}) + \alpha \max\{p_s(x_n, \hat{x}), p_s(x_n, T(\hat{x})), p_s(\hat{x}, T(x_n))\}] \\ &\quad + \frac{\beta}{2} \min\{p_s(x_n, T(\hat{x})), p_s(\hat{x}, T(x_n)), p_s(x_n, T(x_n)) + p_s(\hat{x}, T(\hat{x}))\}] \\ &\leq s[p_s(\hat{x}, x_{n+1}) + \alpha \max\{p_s(x_n, \hat{x}), p_s(x_n, T(\hat{x})), p_s(\hat{x}, T(x_n))\}] \\ &\quad + \frac{\beta}{4}(p_s(x_n, T(\hat{x})) + p_s(\hat{x}, T(x_n)) + p_s(x_n, T(x_n)) + p_s(\hat{x}, T(\hat{x}))). \end{aligned} \quad (9)$$

**Case I:** Suppose  $\max\{p_s(x_n, \hat{x}), p_s(x_n, T(\hat{x})), p_s(\hat{x}, T(x_n))\} = p_s(x_n, \hat{x})$ . Then, from inequality (9), we have

$$\begin{aligned} p_s(\hat{x}, T(\hat{x})) &\leq s[p_s(\hat{x}, x_{n+1}) + \alpha p_s(x_n, \hat{x})] \\ &\quad + \frac{\beta}{2} \left( \frac{p_s(x_n, T(\hat{x})) + p_s(\hat{x}, T(x_n)) + p_s(x_n, T(x_n)) + p_s(\hat{x}, T(\hat{x}))}{2} \right) \\ &\leq s p_s(\hat{x}, x_{n+1}) + s \alpha p_s(x_n, \hat{x}) + \frac{s\beta}{4}(s(p_s(x_n, \hat{x}) + p_s(\hat{x}, T(\hat{x})))) \\ &\quad + p_s(\hat{x}, x_{n+1}) + p_s(x_n, x_{n+1}) + p_s(\hat{x}, T(\hat{x})) \\ &= (s + \frac{s\beta}{4}) p_s(\hat{x}, x_{n+1}) + (s\alpha + \frac{s^2\beta}{4}) p_s(x_n, \hat{x}) \\ &\quad + (\frac{s^2\beta}{4} + \frac{s\beta}{4}) p_s(\hat{x}, T(\hat{x})) + \frac{s\beta}{4} p_s(x_n, x_{n+1}). \end{aligned}$$



From the above inequality, we have

$$(1 - \frac{s^2\beta}{4} - \frac{s\beta}{4})p_s(\hat{x}, T(\hat{x})) \leq (s + \frac{s\beta}{4})p_s(\hat{x}, x_{n+1}) + \\ (s\alpha + \frac{s^2\beta}{4})p_s(x_n, \hat{x}) + \frac{s\beta}{4}p_s(x_n, x_{n+1}),$$

which implies

$$p_s(\hat{x}, T(\hat{x})) \leq \frac{4}{4 - s^2\beta - s\beta} \left[ (s + \frac{s\beta}{4})p_s(\hat{x}, x_{n+1}) + \right. \\ \left. (s\alpha + \frac{s^2\beta}{4})p_s(x_n, \hat{x}) + \frac{s\beta}{4}p_s(x_n, x_{n+1}) \right]. \quad (10)$$

We can observe that for  $\beta \in \min\{\frac{1}{s^3}, \frac{2}{s+1}\}$ ,

$$4 - s^2\beta - s\beta = 4 - s^2\beta - s\beta \\ = 4 - s\beta(s + 1). \quad (11)$$

If  $\beta = \frac{1}{s^3}$ , then, from equality (11) we have

$$4 - s^2\beta - s\beta = 4 - s\beta(s + 1) \\ = 4 - s(s + 1)\frac{1}{s^3} \\ = 4 - \frac{s + 1}{s^2} \\ > 0 \text{ for all } s \geq 1. \quad (12)$$

Similarly, if  $\beta = \frac{2}{s+1}$ , then, from equality (11),

$$4 - s^2\beta - s\beta = 4 - s\beta(s + 1) \\ = 4 - (s + 1)s\frac{2}{s + 1} \\ \leq 4 - s(s + 1)\frac{1}{s^3} \\ > 0 \text{ for all } s \geq 1. \quad (13)$$

From equalities (12) and (13), we conclude that the right-hand side of (10) is non-negative.

**Case II:** Suppose  $\max\{p_s(x_n, \hat{x}), p_s(x_n, T(\hat{x})), p_s(\hat{x}, T(x_n))\} = p_s(x_n, T(\hat{x}))$ . Then, from inequality (9), we have

$$\begin{aligned}
 p_s(\hat{x}, T(\hat{x})) &\leq s[p_s(\hat{x}, x_{n+1}) + \alpha p_s(x_n, T(\hat{x})) \\
 &+ \frac{\beta}{2} \left( \frac{p_s(x_n, T(\hat{x})) + p_s(\hat{x}, T(x_n)) + p_s(x_n, T(x_n)) + p_s(\hat{x}, T(\hat{x}))}{2} \right)] \\
 &\leq s[p_s(\hat{x}, x_{n+1}) + \alpha s(p_s(x_n, \hat{x}) + p_s(\hat{x}, T(\hat{x}))) \\
 &+ \frac{\beta}{4} (s(p_s(x_n, \hat{x}) + p_s(\hat{x}, T(\hat{x}))) + p_s(\hat{x}, x_{n+1}) + p_s(x_n, x_{n+1}) + p_s(\hat{x}, T(\hat{x})))] \\
 &\leq (s + \frac{\beta}{4}s)p_s(\hat{x}, x_{n+1}) + (s^2\alpha + \frac{\beta}{4}s^2)p_s(x_n, \hat{x}) \\
 &+ (s^2\alpha + \frac{\beta}{4}s^2 + \frac{\beta}{4}s)p_s(\hat{x}, T(\hat{x})) + \frac{\beta}{4}s p_s(x_n, x_{n+1}),
 \end{aligned}$$

from the above inequality, we have

$$\begin{aligned}
 (1 - s^2\alpha - \frac{\beta}{4}s^2 - \frac{\beta}{4}s)p_s(\hat{x}, T(\hat{x})) &\leq (s + \frac{\beta}{4}s)p_s(\hat{x}, x_{n+1}) \\
 &+ (s^2\alpha + \frac{\beta}{4}s^2)p_s(x_n, \hat{x}) + \frac{\beta}{4}s p_s(x_n, x_{n+1}),
 \end{aligned}$$

so that

$$\begin{aligned}
 p_s(\hat{x}, T(\hat{x})) &\leq \frac{4}{4 - 4s^2\alpha - \beta s^2 - \beta s} \left[ (s + \frac{\beta}{4}s)p_s(\hat{x}, x_{n+1}) \right. \\
 &\quad \left. + (s^2\alpha + \frac{\beta}{4}s^2)p_s(x_n, \hat{x}) + \frac{\beta}{4}s p_s(x_n, x_{n+1}) \right] \tag{14}
 \end{aligned}$$

from the fact that  $\theta \in \min\{\frac{1}{s^3}, \frac{2}{s+1}\}$ , we have if  $\alpha > \beta$  then by (14), we have

$$\begin{aligned}
 4 - 4s^2\alpha - \beta s^2 - \beta s &= 4 - 4s^2\beta + \beta s^2 + \beta s \\
 &= 4 - (5s + 1)s\beta. \tag{15}
 \end{aligned}$$

If  $\beta = \frac{1}{s^3}$  by (15), we have

$$\begin{aligned}
 4 - 4s^2\alpha - \beta s^2 - \beta s &= 4 - (5s + 1)s\beta \\
 &= 4 - (5s + 1)s\frac{1}{s^3} \\
 &\geq 0 \quad \text{for all } s \geq 1. \tag{16}
 \end{aligned}$$



If  $\beta = \frac{2}{s+1}$  by (15), we have

$$\begin{aligned}
 4 - 4s^2\alpha - \beta s^2 - \beta s &= 4 - (5s + 1)s\beta \\
 &= 4 - (5s + 1)s \frac{2}{s+1} \\
 &\leq 4 - (5s + 1)s \frac{1}{s^3} \\
 &\geq 0 \quad \text{for all } s \geq 1.
 \end{aligned} \tag{17}$$

From inequalities (16) and (17), we conclude that the right-hand side of (10) is non-negative.

If  $\alpha < \beta$ , then by (14), we have

$$\begin{aligned}
 4 - 4s^2\alpha - \beta s^2 - \beta s &= 4 - 4s^2\alpha + \alpha s^2 + \beta s \\
 &= 4 - 5s^2\alpha + s\alpha \\
 &= 4 - (5s + 1)s\alpha.
 \end{aligned} \tag{18}$$

Similarly for (18), we conclude that the right-hand side of (10) is non-negative.

**Case III:** Suppose  $\max\{p_s(x_n, \hat{x}), p_s(x_n, T(\hat{x})), p_s(\hat{x}, T(x_n))\} = p_s(\hat{x}, T(x_n))$ . Then, from inequality (9), we have

$$\begin{aligned}
 p_s(\hat{x}, T(\hat{x})) &\leq s[p_s(\hat{x}, x_{n+1}) + \alpha p_s(\hat{x}, T(x_n))] \\
 &\quad + \frac{\beta}{4}(p_s(x_n, T(\hat{x})) + p_s(\hat{x}, T(x_n)) + p_s(x_n, T(x_n)) + p_s(\hat{x}, T(\hat{x}))) \\
 &\leq s[p_s(\hat{x}, x_{n+1}) + \alpha s p_s(\hat{x}, x_{n+1})] \\
 &\quad + \frac{\beta}{4}(s(p_s(x_n, \hat{x}) + p_s(\hat{x}, T(\hat{x}))) + p_s(\hat{x}, x_{n+1}) + p_s(x_n, x_{n+1}) + p_s(\hat{x}, T(\hat{x}))) \\
 &\leq (s + \alpha s + \frac{\beta}{4}s)p_s(\hat{x}, x_{n+1}) + \frac{\beta}{4}s^2 p_s(x_n, \hat{x}) \\
 &\quad + (\frac{\beta}{4}s^2 + \frac{\beta}{4}s)p_s(\hat{x}, T(\hat{x})) + \frac{\beta}{4}s p_s(x_n, x_{n+1}).
 \end{aligned}$$

By the simplification of the above equality, we have

$$\begin{aligned}
 p_s(\hat{x}, T(\hat{x})) &\leq \frac{4}{4 - s^2\beta - s\beta} [(s + \alpha s + \frac{\beta}{4}s)p_s(\hat{x}, x_{n+1}) \\
 &\quad + \frac{\beta}{4}s^2 p_s(x_n, \hat{x}) + \frac{\beta}{4}s p_s(x_n, x_{n+1})].
 \end{aligned} \tag{19}$$

Note that, for any value of  $\alpha, \beta \in [0, \theta]$  and  $4 - s^2\beta - s\beta \geq 0$ . Thus, the right-hand side of (10) is non-negative. Taking the limit as  $n \rightarrow +\infty$  of both sides in the respective inequalities (10), (14) and (19), we conclude that

$$\begin{aligned} p_s(\hat{x}, T(\hat{x})) &= \lim_{n \rightarrow +\infty} p_s(\hat{x}, T(\hat{x})) \\ &= 0. \end{aligned}$$

Thus,  $T(\hat{x}) = \hat{x}$ .

Next, we prove that if  $\hat{x} \in X$  is a fixed point of  $T$ , then  $p_s(\hat{x}, \hat{x}) = 0$ .

Suppose  $p_s(\hat{x}, \hat{x}) \neq 0$ . Then

$$\begin{aligned} p_s(\hat{x}, \hat{x}) &= p_s(T(\hat{x}), \hat{x}) \\ &\leq \alpha \max\{p_s(\hat{x}, \hat{x}), p_s(\hat{x}, \hat{x}), p_s(\hat{x}, \hat{x})\} \\ &+ \frac{\beta}{2} \min\{p_s(\hat{x}, \hat{x}) + p_s(\hat{x}, T(\hat{x})), p_s(\hat{x}, \hat{x}) + p_s(\hat{x}, T(\hat{x}))\} \\ &= \alpha \max\{p_s(\hat{x}, \hat{x}), p_s(\hat{x}, \hat{x}), p_s(\hat{x}, \hat{x})\} \\ &+ \frac{\beta}{2} \min\{p_s(\hat{x}, \hat{x}) + p_s(\hat{x}, \hat{x}), p_s(\hat{x}, \hat{x}) + p_s(\hat{x}, \hat{x})\} \\ &= (\alpha + \beta)p_s(\hat{x}, \hat{x}) \\ &= \theta p_s(\hat{x}, \hat{x}) \\ &< p_s(\hat{x}, \hat{x}). \end{aligned}$$

This is contradicting the fact that  $p_s(\hat{x}, \hat{x}) \neq 0$ . Therefore,  $p_s(\hat{x}, \hat{x}) = 0$ .

Last, we will prove the uniqueness of the fixed point. Let  $x_1, x_2 \in X$  be two distinct fixed points of  $T$ . Then

$$\begin{aligned} p_s(x_1, x_2) &= p_s(T(x_1), T(x_2)) \\ &\leq \alpha \max\{p_s(x_1, x_2), p_s(x_1, T(x_2)), p_s(x_2, T(x_1))\} \\ &+ \frac{\beta}{2} \min\{p_s(x_1, T(x_2)) + p_s(x_2, T(x_1)), p_s(x_1, T(x_1)) + p_s(x_2, T(x_2))\} \\ &= \alpha \max\{p_s(x_1, x_2), p_s(x_1, x_2), p_s(x_1, x_2)\} \\ &+ \frac{\beta}{2} \min\{p_s(x_1, x_2) + p_s(x_1, x_2), p_s(x_1, x_2) + p_s(x_1, x_2)\} \\ &= (\alpha + \beta)p_s(x_1, x_2) \\ &= \theta p_s(x_1, x_2) \\ &< p_s(x_1, x_2). \end{aligned}$$



This is a contradiction. Therefore, the fixed point is unique.  $\square$

**REMARK 2.** If we take  $\alpha = \frac{\beta}{2}$  and  $p_s(x, T(y)) + p_s(y, T(x)) \geq p_s(x, T(x)) + p_s(y, T(y))$  then we find Theorem 1 of Batsari et al. ([Batsari & Kumam, 2020](#)).

**COROLLARY 3.** Let  $(X, p)$  be a complete partial metric space associated with a partial order  $\preceq$ . Suppose an order-preserving mapping  $T : X \rightarrow X$  satisfies

$$\begin{aligned} p_s(T(x), T(y)) &\leq \alpha \max\{p_s(x, y), p_s(x, T(y)), p_s(y, T(x))\} \\ &+ \frac{\beta}{2} \min\{p_s(x, T(y)) + p_s(y, T(x)), p_s(x, T(x)) + p_s(y, T(y))\} \end{aligned} \quad (20)$$

for all comparable  $x, y \in X$ , where  $\theta \in [0, 1]$ . If there exists  $x_0 \in X$  such that  $x_0 \preceq T(x_0)$ , then  $T$  has a unique fixed point  $\hat{x} \in X$  and  $p(\hat{x}, \hat{x}) = 0$ .

Now we apply our main result similar to ([Batsari & Kumam, 2020](#)) as follows:

### Application to quantum operations

In quantum systems, measurements can be seen as quantum operations ([Seevinck, 2003](#)). Quantum operations are very important in narrating quantum systems that collaborate with the environment.

Let  $\mathcal{B}(H)$  be the set of bounded linear operators on the separable complex Hilbert space  $H$ ;  $\mathcal{B}(H)$  is the state space of consideration. Suppose  $\mathcal{A} = \{A_i, A_i^* : i = 1, 2, 3, \dots\}$  is a collection of operators  $A'_i \in \mathcal{B}(H)$  satisfying  $\sum A_i A_i^* \leq I$ . A map  $\phi : \mathcal{B}(H) \rightarrow \mathcal{B}(H)$  of the form  $\phi_{\mathcal{A}}(B) = \sum A_i B A_i^*$  is called a quantum operation ([Arias et al, 2002](#)), quantum operations can be used in quantum measurements of states. If the  $A_i$ 's are self adjoint then,  $\phi_{\mathcal{A}}$  is self-adjoint.

General quantum measurements that have more than two values are narrated by effect-valued measures ([Arias et al, 2002](#)). Denote the set of quantum effects by  $\varepsilon(H) = \{A \in \mathcal{B}(H) : 0 \leq A \leq I\}$ . Consider the discrete effect-valued measures narrated by a sequence of  $E_i \in \varepsilon(H)$ ,  $i = 1, 2, \dots$  satisfying  $E_i = I$  where the sum converges in the strong operator topology. Therefore, the probability that outcome  $i$  eventuates in the state  $\rho$  is  ${}_{\rho}(E_i)$

and the post-measurement state given that  $i$  eventuates is  $\frac{E_i^{\frac{1}{2}} \rho E_i^{\frac{1}{2}}}{tr \rho E_i}$  (Arias et al, 2002). Furthermore, the resulting state after the implementation of measurement without making any consideration is given by

$$\phi(\rho) = \sum E_i^{\frac{1}{2}} \rho E_i^{\frac{1}{2}} \quad (21)$$

If the measurement does not disturb the state  $\rho$ , then we have  $\phi(\rho) = \rho$ . Furthermore, the probability that an effect  $A$  eventuates in the state  $r$  given that the measurement was conducted is

$$P_{\phi(\rho)}(A) = tr \left[ A \sum E_i^{\frac{1}{2}} \rho E_i^{\frac{1}{2}} \right] = tr \left( \sum E_i^{\frac{1}{2}} \rho E_i^{\frac{1}{2}} \rho \right) \quad (22)$$

If  $A$  is not interrupted by the measurement in any state we have

$$\sum E_i^{\frac{1}{2}} \rho E_i^{\frac{1}{2}} = A,$$

and by defining  $\phi(A) = \sum E_i^{\frac{1}{2}} \rho E_i^{\frac{1}{2}}$ , we end up with  $\phi(A) = A$ .

From now, we will be dealing with a bi-level ( $|0\rangle, |1\rangle$ ) single qubit quantum system where a quantum state  $|\Psi\rangle$  can be narrated as

$$|\Psi\rangle = a|0\rangle + b|1\rangle, \text{ with } a, b \in \mathbb{C} \text{ and } |a|^2 + |b|^2 = 1$$

see (Batsari & Kumam, 2020; Nielsen & Chuang, 2000). Considering the characterization of a bi-level quantum system by the Bloch sphere (Figure 1) above, a quantum state ( $|\Psi\rangle$ ) can be represented with the density matrix below ( $\rho$ ),

$$|\Psi\rangle = \rho = \frac{1}{2} \begin{pmatrix} 1 + \eta \cos \theta & \eta \cos \phi \sin \theta - i \eta \sin \phi \sin \theta \\ \eta \cos \phi \sin \theta + i \eta \sin \phi \sin \theta & 1 - \eta \cos \theta \end{pmatrix},$$

$\eta \in [0, 1], 0 \leq \theta \leq \pi, \text{ and } 0 \leq \phi \leq 2\pi.$       (23)

Also, the density ( $\rho$ ) matrix is,

$$\rho = \frac{1}{2} [I + \bar{r}_\rho \cdot \bar{\sigma}] = \frac{1}{2} \begin{bmatrix} 1 + r_z & r_x - ir_y \\ r_x + ir_y & 1 - r_z \end{bmatrix} \quad (24)$$

where  $\bar{r}_\rho = [r_x, r_y, r_z]$  is the Bloch vector with  $\|\bar{r}_\rho\| \leq 1$ , and  $\bar{\sigma} = [\sigma_x, \sigma_y, \sigma_z]$  where

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$




Figure 1 – Bloch sphere

Рис. 1 – Сфера Блоха

Слика 1 – Блохова сфера

Let  $\rho, \sigma$  be two quantum states in a bi-level quantum system. Then, the Bures fidelity (Bures, 1969) between the quantum states  $\rho$  and  $\sigma$  is defined as

$$F(\rho, \sigma) = [tr\sqrt{\rho^{\frac{1}{2}}\sigma\rho^{\frac{1}{2}}}]^2.$$

The Bures fidelity satisfies  $0 \leq F(\rho, \sigma) \leq 1$ , if  $\rho = \sigma$  it takes the value 1 and 0 if  $\rho$  and  $\sigma$  have an orthogonal support (Nielsen & Chuang, 2000).

Now consider a two-level quantum system  $X$  represented with the collection of density matrices  $\{\rho : \rho \text{ is as defined in Equation (24)}\}$ . Define the function  $p_s : X \times X \rightarrow \mathbb{R}_+$  as follows:

$$p_s(\rho, \delta) = \begin{cases} \max\{\|\overline{r_\rho}\|, \|\overline{r_\delta}\|\} e^{\frac{1}{6}(1-F(\rho, \delta))}, & \rho \neq \delta, \\ 0, & \rho = \delta. \end{cases}$$

It is easy to show that  $p_s$  is a b-metric on  $X$  with  $s$  taking the value 1 approximately. They also define an order relation  $\preceq$  on  $X$  by

$\rho \preceq \delta$  iff the line from the origin joining the point  $\overline{r_\delta}$  passes through  $\overline{r_\rho}$ .  
(25)

It is easy to show that the order relation defined above is a partial order ([Batsari & Kumam, 2020](#)).

As in ([Batsari & Kumam, 2020](#)), we find the following corollary.

**COROLLARY 4.** Let  $(p_s, X)$  be a complete partial b-metric space associated with the above order  $\preceq$ . Suppose an order-preserving quantum operation  $T : X \rightarrow X$  that satisfies conditions in Theorems 1. Then,  $T$  has a fixed point.

The following example validates our main result.

**EXAMPLE 0.1.** Consider the depolarizing quantum operation  $T$  on the Bloch sphere  $X$ ;  $T(\rho) = \frac{I}{2}p + (1-p)\rho$  with the depolarizing parameter  $p \in [0, 1]$ . Let the comparable quantum states satisfy (25).

We examine that  $T : X \rightarrow X$  satisfies all the conditions of our theorem. Now, let  $\rho, \delta \in X$ . We show that  $T$  is order preserving with definition (25). For this, we will prove that if  $\rho \preceq \delta$  then  $T(\rho) \preceq T(\delta)$ .

Therefore, as ([Batsari & Kumam, 2020](#)) using the Bloch sphere representation of states in a bi-level quantum system below

$$\rho = \frac{1}{2} \begin{pmatrix} 1 + \mu \cos \theta & \mu \cos \phi \sin \theta - i\mu \sin \phi \sin \theta \\ \mu \cos \phi \sin \theta + i\mu \sin \phi \sin \theta & 1 - \mu \cos \theta \end{pmatrix},$$

$\mu \in [0, 1], 0 \leq \theta \leq \pi, \text{ and } 0 \leq \phi \leq 2\pi,$

So,

$$\begin{aligned} T(\rho) &= \frac{1}{2} \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} + \\ &\quad \frac{1-p}{2} \begin{pmatrix} 1 + \mu \cos \theta & \mu \cos \phi \sin \theta - i\mu \sin \phi \sin \theta \\ \mu \cos \phi \sin \theta + i\mu \sin \phi \sin \theta & 1 - \mu \cos \theta \end{pmatrix} \\ &= \frac{1}{2} \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 1 + \mu \cos \theta & \mu \cos \phi \sin \theta - i\mu \sin \phi \sin \theta \\ \mu \cos \phi \sin \theta + i\mu \sin \phi \sin \theta & 1 - \mu \cos \theta \end{pmatrix} \\ &\quad - \frac{1}{2} \begin{pmatrix} p + p\mu \cos \theta & p\mu \cos \phi \sin \theta - ip\mu \sin \phi \sin \theta \\ p\mu \cos \phi \sin \theta + ip\mu \sin \phi \sin \theta & p - p\mu \cos \theta \end{pmatrix} \\ &= \frac{1}{2} \begin{pmatrix} 1+(1-p)\mu \cos \theta & (1-p)[\mu \cos \phi \sin \theta - i\mu \sin \phi \sin \theta] \\ (1-p)[\mu \cos \phi \sin \theta + i\mu \sin \phi \sin \theta] & 1-(1-p)\mu \cos \theta \end{pmatrix} \end{aligned}$$



Clearly, the angles  $\theta$  and  $\phi$  do not change by the depolarizing quantum operation  $T$ . Also, we can deduce that the distance of the quantum state  $\rho$  from the origin given by  $\mu$  is greater than or equal to the distance of the new quantum state  $T(\rho)$  from the origin given by  $(1-p)\mu$ ,  $p \in [0, 1]$ . Consequently, for any two quantum states which are comparable  $\rho, \delta \in X$  ( $\rho \preceq \delta$ ), with respective distances from the origin  $\mu_\rho$  and  $\mu_\delta$  such that,  $\mu_\rho \leq \mu_\delta$ , the depolarizing quantum operation  $T$  constructs two quantum states  $T(\rho), T(\delta) \in X$ , have distances  $(1-p)\mu_\rho$  and  $(1-p)\mu_\delta$  from the origin for  $p \in [0, 1]$  respectively. Since  $\mu_\rho \leq \mu_\delta$ , then  $(1-p)\mu_\rho \leq (1-p)\mu_\delta$ , for all  $p \in [0, 1]$ . Thus,  $T(\rho) \preceq T(\delta)$ , which proves  $T$  is order-preserving.

The fidelity of any two quantum states  $\rho = \frac{1}{2}(I_2 + \vec{r}_\rho \cdot \vec{\sigma})$  and  $\delta = \frac{1}{2}(I_2 + \vec{r}_\delta \cdot \vec{\sigma})$  is,

$$F(\rho, \delta) = \frac{1}{2}[1 + \vec{r}_\rho \cdot \vec{r}_\delta + \sqrt{1 - \|\vec{r}_\rho\|^2} \sqrt{1 - \|\vec{r}_\delta\|^2}] \quad (26)$$

see (Batsari & Kumam, 2020; Chen et al, 2002), where  $\vec{r}_\rho \cdot \vec{r}_\delta$  is the inner dot product between the vectors  $\vec{r}_\rho$  and  $\vec{r}_\delta$ . So, for any comparable quantum states  $\rho = \frac{1}{2}(I_2 + \vec{r}_\rho \cdot \vec{\sigma})$  and  $\delta = \frac{1}{2}(I_2 + \vec{r}_\delta \cdot \vec{\sigma})$ ,  $\vec{r}_\rho \cdot \vec{r}_\delta = \|\vec{r}_\rho\| \|\vec{r}_\delta\| \cos \vartheta$  for  $\vartheta$  being the angle between  $\vec{r}_\rho$  and  $\vec{r}_\delta$ . Using Equation (26), we have,

(i).  $F(\rho, \rho) = 1$ .

(ii)  $F(\rho, o) = \frac{1}{2}$ ; for  $\rho$  a pure state and  $o$  the completely mixed state.

(iii)  $F(\rho, \rho-) = 0$ ; for  $\rho$  a pure state that is  $180^\circ$  separated from  $\rho$ ,

see (Davies, 1976; Göhde, 1965). Thus,  $1.000 \leq e^{\frac{1}{6}(1-F(\rho,\delta))} \leq 1.181$  for  $\rho, \delta \in X$ . Now, using  $s = 1$  and  $\theta \in [0, 1]$  on Theorems 1. We have

$$\begin{aligned} p_s(T(\rho), T(\delta)) &= \max\{\|\vec{r}_\rho\|, \|\vec{r}_\delta\|\} e^{\frac{1}{5}(1-F(T(\rho), T(\delta)))} \\ &= \frac{1}{4} \|\delta\| e^{\frac{1}{5}(1-F(T(\rho), T(\delta)))} \\ &\leq \frac{1}{4} (\|\delta\| e^{\frac{1}{5}(1-F(T(\rho), \delta))} + \|\rho\| e^{\frac{1}{5}(1-F(T(\rho), \rho))}) \\ &= \frac{1}{2} \left( \frac{1}{2} (p_s(T(\rho), \delta) + p_s(T(\rho), \rho)) \right) \\ &= \frac{1}{2} \left( \frac{1}{2} \max\{p_s(\rho, \delta), p_s(\rho, T(\delta)), p_s(T(\rho), \delta)\} \right. \\ &\quad \left. + \min\left\{\frac{p_s(\rho, T(\delta)) + p_s(T(\rho), \delta)}{2}, \frac{p_s(\rho, T(\rho)) + p_s(\delta, T(\delta))}{2}\right\} \right). \end{aligned}$$

Taking  $\alpha = \frac{1}{2}$  and  $\beta = 1$ , condition (1) in Theorem 1 is satisfied. So  $T$  has a unique fixed point in  $X$ .

## References

- Aamri, M. & El Moutawakil, D. 2002. Some new common fixed point theorems under strict contractive conditions. *Journal of Mathematical Analysis and Applications*, 270(1), pp.181-188. Available at: [https://doi.org/10.1016/S0022-247X\(02\)00059-8](https://doi.org/10.1016/S0022-247X(02)00059-8).
- Agarwal, R.P., Karapinar, E., O'Regan, D. & Roldán-López-de-Hierro, A.F. 2015. *Fixed point theory in metric type spaces*. Springer, Cham. Available at: <https://doi.org/10.1007/978-3-319-24082-4>. ISBN: 978-3-319-24082-4.
- Arias, A., Gheondea, A. & Gudder, S. 2002. Fixed Points of Quantum Operations. *Journal of Mathematical Physics*, 43(12), pp.5872-5881. Available at: <https://doi.org/10.1063/1.1519669>.
- Bakhtin, I. 1989. The contraction mapping principle in quasimetric spaces. *Func. An., Gos. Ped. Inst. Unianowsk*, 30, pp.26-37.
- Banach, S. 1922. Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales. *Fundamenta Mathematicae*, 3, pp.133-181 (in French). Available at: <https://doi.org/10.4064/fm-3-1-133-181>.
- Batsari, U.Y. & Kumam, P. 2018. A Globally Stable Fixed Point in an Ordered Partial Metric Space. In: Anh, L., Dong, L., Kreinovich, V. & Thach, N. (Eds.) *Econometrics for Financial Applications. ECONVN 2018. Studies in Computational Intelligence*, 760, pp.360-368. Springer, Cham. Available at: [https://doi.org/10.1007/978-3-319-73150-6\\_29](https://doi.org/10.1007/978-3-319-73150-6_29).
- Batsari, U.Y. & Kumam, P. 2020. Some Generalised Fixed Point Theorems Applied to Quantum Operations. *Symmetry*, 12(5), art.ID:759. Available at: <https://doi.org/10.3390/sym12050759>.
- Batsari, U.Y., Kumam, P. & Sitthithakerngkiet, K. 2018. Some globally stable fixed points in b-metric spaces. *Symmetry*, 10(11), art.ID:555. Available at: <https://doi.org/10.3390/sym10110555>.
- Bourbaki, N. 1974. *Topologie Generale*. Paris, France: Herman. ISBN-13: 978-2705656928.
- Brouwer, L.E.J. 1911. Über Abbildung von Mannigfaltigkeiten. *Mathematische Annalen*, 71, pp.97-115. Available at: <https://doi.org/10.1007/BF01456931>.
- Browder, F.E. 1959. On a generalization of the Schauder fixed point theorem. *Duke Mathematical Journal*, 26(2), pp.291-303. Available at: <https://doi.org/10.1215/S0012-7094-59-02629-8>.
- Bures, D. 1969. An Extension of Kakutani's Theorem on Infinite Product Measures to the Tensor Product of Semifinite  $w^*$ -Algebras. *Transactions of the American Mathematical Society*, 135, pp.199-212. Available at: <https://doi.org/10.2307/1995012>.
- Busch, P. & Singh, J. 1998. Lüders Theorem for Unsharp Quantum Measurements. *Physics Letters A*, 249(1-2), pp.10-12. Available at: [https://doi.org/10.1016/S0375-9601\(98\)00704-X](https://doi.org/10.1016/S0375-9601(98)00704-X).



- Chen, J-L., Fu, L., Ungar, A.A. & Zhao, X-G. 2002. Alternative fidelity measure between two states of an  $N$ -state quantum system. *Physical Review A*, 65(art.number:054304). Available at: <https://doi.org/10.1103/PhysRevA.65.054304>.
- Chidume, C.E. & Chidume, C.O. 2014. *Foundations of Mathematical Analysis*. Ibadan, Nigeria: University of Ibadan, Ibadan University Press Publishing House. ISBN: 978-978-8456-32-2.
- Czerwinski, S. 1993. Contraction mappings in b-metric spaces. *Acta Mathematica et Informatica Universitatis Ostraviensis*, 1(1), pp.5-11 [online]. Available at: <https://dml.cz/handle/10338.dmlcz/120469> [Accessed: 20 March 2022].
- Davies, E.B. 1976. *Quantum Theory of Open Systems*. London, UK: Academic Press. ISBN-13: 978-0122061509.
- Debnath, P., Konwar, N. & Radenović, S. 2021. *Metric Fixed Point Theory: Applications in Science, Engineering and Behavioural Sciences*. Springer Verlag, Singapore. Available at: <https://doi.org/10.1007/978-981-16-4896-0>.
- Du, W-S., Karapinar, E. & He, Z. 2018. Some Simultaneous Generalizations of Well-Known Fixed Point Theorems and Their Applications to Fixed Point Theory. *Mathematics*, 6(7), art.ID:117. Available at: <https://doi.org/10.3390/math6070117>.
- Göhde, D. 1965. Zum Prinzip der Kontraktiven abbildung. *Mathematische Nachrichten*, 30(3-4), pp.251-258. Available at: <https://doi.org/10.1002/mana.19650300312>.
- Kannan, R. 1972. Some results on fixed points - IV. *Fundamenta Mathematicae*, 74, pp.181-187. Available at: <https://doi.org/10.4064/fm-74-3-181-187>.
- Khan, M.S., Swaleh, M. & Sessa, S. 1984. Fixed point theorems by altering distances between the points. *Bulletin of the Australian Mathematical Society*, 30(1), pp.1-9. Available at: <https://doi.org/10.1017/S0004972700001659>.
- Kirk, W. & Shahzad, N. 2014. *Fixed Point Theory in Distance Spaces*. Springer International Publishing Switzerland. Available at: <https://doi.org/10.1007/978-3-319-10927-5>.
- Knaster, B. 1928. Un theoreme sur les fonctions denses. *Annales de la Société polonaise de mathématique*, 6, pp.133-134.
- Leray, J. & Schauder, J. 1934. Topologie et équations fonctionnelles. *Annales scientifiques de l'École Normale Supérieure*, 51, pp.45-78. Available at: <https://doi.org/10.24033/asens.836>.
- Long, L. & Zhang, S. 2011. Fixed points of commutative super-operators. *Journal of Physics A: Mathematical and Theoretical*, 44(9), art.ID:095201. Available at: <https://doi.org/10.1088/1751-8113/44/9/095201>.
- Lüders, G. 1950. Über die Zustandsänderung durch den Meßprozeß. *Annalen der physik*, 443(5-8), pp.322-328. Available at: <https://doi.org/10.1002/andp.19504430510>.

- Matthews, S.G. 1994. Partial Metric Topology. *Annals of the New York Academy of Sciences*, 728(1) General Topology and Applications, pp.183-197. Available at: <https://doi.org/10.1111/j.1749-6632.1994.tb44144.x>.
- Nielsen, M.A. & Chuang, I.L. 2000. *Quantum Computation and Quantum Information*. Cambridge, UK: Cambridge University Press. Available at: <https://doi.org/10.1017/CBO9780511976667>.
- Schauder, J. 1930. Der Fixpunktsatz in Funktionalräumen. *Studia Mathematica*, 2(1), pp.171-180 [online]. Available at: <https://eudml.org/doc/urn:eudml:doc:217247> [Accessed: 22 March 2022].
- Seevinck, M.P. 2003. *Quantum Operations and Measurement*, 2nd ed. Utrecht, The Netherlands: Utrecht University [online]. Available at: <http://mpseevinck.ruhousing.nl/seevinck/lezingoviedo03a.pdf> [Accessed: 22 March 2022].
- Shukla S. 2014. Partial b-metric spaces and fixed point theorems. *Mediterranean Journal of Mathematics*, 11, pp.703-711. Available at: <https://doi.org/10.1007/s00009-013-0327-4>.
- Tarski, A. 1955. A lattice-theoretical fixpoint theorem and its applications. *Pacific Journal of Mathematics*, 5(2), pp.285-309. Available at: <https://doi.org/10.2140/pjm.1955.5.285>.
- Tarski, A.A. 1949. A fixpoint theorem for lattices and its applications (preliminary report). *Bulletin of the American Mathematical Society*, 1949(55), 1051-1052.
- Zhang, H. & Ji, G. 2012. A Note on Fixed Points of General Quantum Operations. *Reports on Mathematical Physics*, 70(1), pp.111-117. Available at: [https://doi.org/10.1016/S0034-4877\(13\)60016-6](https://doi.org/10.1016/S0034-4877(13)60016-6).
- Zhang, H. & Si, H. 2016. Fixed Points Associated to Power of Normal Completely Positive Maps\*. *Journal of Applied Mathematics and Physics*, 4(5), pp.925-929. Available at: <https://doi.org/10.4236/jamp.2016.45101>.

---

ТЕОРЕМА О ФИКСИРОВАННОЙ ТОЧКЕ В ЧАСТИЧНОМ  
b-МЕТРИЧЕСКОМ ПРОСТРАНСТВЕ С ПРИМЕНЕНИЕМ В  
КВАНТОВЫХ ОПЕРАЦИЯХ

Ракеш Тивари<sup>a</sup>, Мухаммад Саид Khan<sup>b</sup>,  
Шоба Рани<sup>a</sup>, Никола Фабиано<sup>b</sup>

<sup>a</sup> Государственный автономный колледж последипломного образования VYT , кафедра математики, г. Дург, Чхаттисгарх, Республика Индия

<sup>b</sup> Университет медицинских наук Сефако Макгато, кафедра математики и прикладной математики, г. Га-Ранкува, Южно-Африканская Республика



<sup>b</sup> Белградский университет, Институт ядерных наук Винча - Национальный институт Республики Сербия, г. Белград, Республика Сербия, **корреспондент**

РУБРИКА ГРНТИ: 27.35.33 Математические модели  
электродинамики и оптики  
27.35.57 Математические модели квантовой  
физики  
27.39.27 Нелинейный функциональный  
анализ

ВИД СТАТЬИ: оригинальная научная статья

*Резюме:*

*Введение/цель: Сконструирована теорема о неподвижной точке с сохранением порядка в полном и частичном  $b$ -метрическом пространстве при выполнении условий сжатия.*

*Методы: В данной статье применен метод расширения результатов Батсари и др.*

*Результаты: Точность квантового состояния используется для построения неподвижного квантового состояния.*

*Выводы: Неподвижное квантовое состояние связано с квантовой операцией, сохраняющей порядок.*

*Ключевые слова: частичное  $b$ -метрическое пространство, отображение с сохранением порядка, квантовая операция, точность квантового состояния, вектор Блоха.*

---

#### ТЕОРЕМА ФИКСНЕ ТАЧКЕ У ДЕЛИМИЧНОМ $b$ -МЕТРИЧКОМ ПРОСТОРУ ПРИМЕЊЕНА НА КВАНТНЕ ОПЕРАЦИЈЕ

Rakesh Тивари<sup>a</sup>, Мохамед Саид Кан<sup>b</sup>,  
Шоба Рани<sup>a</sup>, Никола Фабиано<sup>Г</sup>

<sup>a</sup> Државни колеџ за последипломске студије В.Ј.Т, Департман за математику, Дург, Чатисгар, Република Индија

<sup>b</sup> Универзитет здравствених наука Сефако Макгато, Департман за математику и примењену математику,  
Га-Ранкува, Јужноафричка Република

<sup>b</sup> Универзитет у Београду, Институт за нуклеарне науке "Винча" - Национални институт Републике Србије, Београд,  
Република Србија, **автор за преписку**

ОБЛАСТ: математика

ВРСТА ЧЛАНКА: оригинални научни рад

**Сажетак:**

**Увод/циљ:** Конструисана је теорема фиксне тачке мапирања с очувањем редоследа на комплетном парцијалном  $b$ -метричком простору уз задовољавање контрактивног услова.

**Методе:** Примењен метод проширен је резултатима Батсарија и других.

**Резултати:** Верност квантног стања користи се за конструисање фиксног квантног стања.

**Закључак:** Фиксно квантно стање повезано је са квантном операцијом која чува редослед.

**Кључне речи:** парцијални  $b$ -метрички простор, мапирање с очувањем редоследа, квантна операција, верност квантног стања, Блохов вектор.

---

Paper received on / Дата получения работы / Датум пријема чланка: 22.03.2022.  
Manuscript corrections submitted on / Дата получения исправленной версии работы /  
Датум достављања исправки рукописа: 21.06.2022.

Paper accepted for publishing on / Дата окончательного согласования работы / Датум коначног прихватања члана за објављивање: 23.06.2022.

© 2022 The Authors. Published by Vojnotehnički glasnik / Military Technical Courier (<http://vtg.mod.gov.rs>, <http://втг.мо.упр.срб>). This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/rs/>).

© 2022 Авторы. Опубликовано в "Военно-технический вестник / Vojnotehnički glasnik / Military Technical Courier" (<http://vtg.mod.gov.rs>, <http://втг.мо.упр.срб>). Данная статья в открытом доступе и распространяется в соответствии с лицензией "Creative Commons" (<http://creativecommons.org/licenses/by/3.0/rs/>).

© 2022 Аутори. Објавио Војнотехнички гласник / Vojnotehnički glasnik / Military Technical Courier (<http://vtg.mod.gov.rs>, <http://втг.мо.упр.срб>). Ово је чланак отвореног приступа и дистрибуира се у складу са Creative Commons лиценцом (<http://creativecommons.org/licenses/by/3.0/rs/>).

